

Additions to Acrocalymmaceae and Didymosphaeriaceae (Pleosporales, Dothideomycetes): Some interesting novel additions from plant litter in China

Danushka S. Tennakoon¹, Nimali I. de Silva¹, Sinang Hongsanan¹, Ning Xie¹

¹ Shenzhen Key Laboratory of Microbial Genetic Engineering, College of Life Science and Oceanography, Shenzhen University, Shenzhen 518060, China
Corresponding author: Ning Xie (ning.xie@szu.edu.cn)



This article is part of:
Exploring the Hidden Fungal Diversity: Biodiversity, Taxonomy, and Phylogeny of Saprobiic Fungi

Edited by Samantha C. Karunarathna,
Danushka Sandaruwan Tennakoon,
Ajay Kumar Gautam

Academic editor: Samantha C.
Karunarathna

Received: 27 June 2025

Accepted: 18 August 2025

Published: 5 September 2025

Citation: Tennakoon DS, de Silva NI, Hongsanan S, Xie N (2025) Additions to Acrocalymmaceae and Didymosphaeriaceae (Pleosporales, Dothideomycetes): Some interesting novel additions from plant litter in China. MycoKeys 122: 59–98. <https://doi.org/10.3897/mycokeys.122.163383>

Copyright: © Danushka S. Tennakoon et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

During a survey of fungal diversity of plant litter in China, numerous isolates with affinity to Pleosporales (Dothideomycetes, Ascomycota) were recovered. Morphology, coupled with phylogenetic analyses using Maximum Likelihood and Bayesian Inference on combined LSU, SSU, ITS, *tub2*, and *tef1*-α sequence data, revealed that these isolates belong to the families Acrocalymmaceae and Didymosphaeriaceae, and are introduced as *Acrocalymma poaceicola* **sp. nov.** and *Neptunomyces chinensis* **sp. nov.** In addition, five new host records, *A. ampeli*, *Neokalmusia arundinis*, *Paraconiothyrium archidendri*, *Pseudopithomyces chartarum*, and *Spegazzinia deightonii* are reported from litter of *Livistona chinensis*, *Sporobolus alterniflorus*, *Citrus maxima*, *Hedychium coronarium*, and *Arundo pliniana*, respectively. All the new species and host records were compared with phylogenetically and morphologically closely related species, and detailed descriptions, micrographs, and phylogenetic analysis results are provided.

Key words: Multi-gene phylogeny, new host records, new species, saprobes, systematics, taxonomy

Introduction

Plant litter, composed of dead plant materials, such as leaves, twigs, bark, seeds, flowers, roots, and other wood debris accumulates naturally on the forest floor (Wardle et al. 2002; Berg and McClaugherty 2008; Liu et al. 2022). Plant litter plays a fundamental role in ecosystem dynamics through the decomposition process (Schroeter et al. 2022), and is crucial for nutrient cycling, improving soil structure, enhancing biodiversity, and boosting forest productivity and resilience (Sayer 2006; Pei et al. 2017; Giweta 2020; Pascoal et al. 2021). The significance of plant litter extends beyond its ecological functions, as it also serves as a habitat for a wide range of organisms, including bacteria, fungi, and macroinvertebrates. Of them, fungal communities are regarded as the “key players” in the decomposition of plant litter, because of their capacity to produce a vast array of extracellular enzymes (Geethanjali and Jayashankar 2016; Asplund et al. 2018; Zhou et al. 2023). They are responsible for the breakdown of plant litter in various ecosystems, significantly contributing to nutrient

cycling and the decomposition of organic matter (Griffiths et al. 2021; Pascoal et al. 2021; Tennakoon et al. 2021; Liu et al. 2022). Hence, it is vital to explore the fungal diversity residing in plant litter to comprehend the relationships involved in plant litter decomposition. Many researchers have conducted studies on fungal species inhabiting plant litter, yielding numerous new findings (Liu et al. 2015; Tanaka et al. 2015; Hyde et al. 2016; Hashimoto et al. 2017; Crous et al. 2020; Tennakoon et al. 2024). Particularly, we are exploring the hidden diversity of Dothideomycetes fungi in plant litter through comprehensive morphology and phylogenetic analyses (Tennakoon et al. 2018, 2020, 2023, 2024; Hyde et al. 2020; Hongsanan et al. 2020; Wanasinghe et al. 2022).

Acrocalymmaeae and Didymosphaeriaceae are two of the speciose families in the Pleosporales (Dothideomycetes, Ascomycota). Of them, Acrocalymmaeae was established by Trakunyingcharoen et al. (2014) to include the monotypic genus *Acrocalymma* with *A. medicaginis* as the type species. Most of the Acrocalymmaeae species have been reported in their asexual morph, characterized by immersed to semi-immersed, globose to sub-globose conidiomata and cylindrical to fusiform, hyaline or pale brown, aseptate, or 1–3-septate conidia with appendages at both ends (Hongsanan et al. 2020; Mortimer et al. 2021; Calabon et al. 2023). The sexual morph has immersed to erumpent, globose to sub-globose, unilocular ascomata, cylindrical asci, and narrowly fusiform, hyaline to brown or pale brown, 1–3-septate ascospores with a mucoid sheath (Jayasiri et al. 2019; Mortimer et al. 2021; Tennakoon et al. 2021; Konta et al. 2023). They have been reported from worldwide, including temperate, subtropical, and tropical regions (e.g., Australia, China, Egypt, India, Spain, Thailand, and the United States) (Shoemaker et al. 1991; Trakunyingcharoen et al. 2014; Dong et al. 2020; Tennakoon et al. 2021; Calabon et al. 2023; Konta et al. 2023). Most species have been found as saprobes on plant litter, which includes dead leaves, wood, pods, and seeds, in both terrestrial and freshwater habitats (Jayasiri et al. 2019; Mortimer et al. 2021; Tennakoon et al. 2021). Some species (e.g., *A. medicaginis* and *A. vagum*) have also been confirmed as plant pathogens (Alcorn and Irwin 1987; Farr et al. 1998; Trakunyingcharoen et al. 2014). Additionally, they have been reported as endophytes, for instance, *A. vagum* from wild rice (Zeng et al. 2023).

Didymosphaeriaceae, introduced by Munk (1956), is typified by the genus *Didymosphaeria*, and *D. epidermidis* is the type species. The morphological characteristics of Didymosphaeriaceae are highly diverse, with most sexual morphs having 1–3-septa (e.g., *Barria*, *Didymosphaeria*, *Kalmusia*, *Letendraea*, and *Neokalmusia*) or muriform (e.g., *Austropleospora*, *Bimuria*, *Deniquelata*, *Julella*, *Karstenula*, and *Tremateia*), brown ascospores and trabeculate pseudoparaphyses that are anastomosed above the asci (Ariyawansa et al. 2014a, b; Liu et al. 2015; Hongsanan et al. 2020). Asexual morphs can be either coelomycetes (e.g., *Alloconiothyrium*, *Austropleospora*, *Letendraea*, *Paraconiothyrium*, and *Paraphaeosphaeria*) or hyphomycetes (e.g., *Pseudopithomyces*, and *Spegazzinia*) (Hyde et al. 2013; Wanasinghe et al. 2016; Hongsanan et al. 2020). The ordinal placement of Didymosphaeriaceae has been a subject of controversy for a long time. Initially, this was placed in Melanommatales based on trabeculate pseudoparaphyses (Barr 1990). Subsequently, Ariyawansa et al. (2014a) provided a comprehensive taxonomic revision of this group, revealing that the family appears to be a distinct family within Pleosporales. Most of the Didymosphaeriaceae species have been recognized as saprobes in terrestrial, freshwater, marine, and mangrove ecosystems (Ariyawansa et al. 2014a, b;

Liu et al. 2015; Phookamsak et al. 2019; Tennakoon et al. 2022). Additionally, some taxa have been reported as both endophytes and pathogens in plants (Yuan et al. 2020; Guarnaccia et al. 2022). Currently, there are 33 accepted genera in this family (Wijayawardene et al. 2022, Htet et al. 2024).

During a mycological survey conducted in China, we isolated seven saprobic fungi from plant litter substrates. The objectives of this study are to identify the isolated taxa using both morphological and phylogenetic approaches. Our findings contribute to the growing body of knowledge on fungal diversity in China.

Materials and methods

Sample collection and examination

Dead wood and leaf specimens were collected from Guangdong and Yunnan Provinces, China (during 2017–2024). The collected specimens were placed in zip-lock bags with collection information (Rathnayaka et al. 2025), brought to the laboratory, and incubated in a plastic box with wet tissue paper (two days). The specimens were initially checked using a stereomicroscope (AXIOSKOP 2 PLUS Series, Göttingen, Germany). All the surface characteristics of ascomata and conidiomata, such as color, shape, and position of the host surface (e.g., immersed, semi-immersed, and erumpent or superficial), were recorded. Hand sections of ascomata and conidiomata were obtained manually using a razor blade and were mounted in distilled water. Micro-morphological characteristics (e.g., ascospores, asci, conidia, conidiogenous cells, peridium, pseudoparaphyses) were examined and photographed using a Nikon ECLIPSE Ni-U compound microscope (Nikon Corporation, Japan), equipped with a Canon AxioCam 506 color digital camera (Hanover, Germany). Lactoglycerol was used to prepare the permanent slides, and nail polish was applied to the margins of the cover slips to seal them. All measurements were calculated using the Tarosoft (R) Image Framework application. Adobe Photoshop CS3 Extended version 10.0 software (Adobe Systems, USA) was used to construct the photo plates.

The approach described by Senanayake et al. (2022) was used for single-spore isolation. After germination, spores were carefully transferred to Potato Dextrose Agar (PDA) and incubated at 25 °C. Pure cultures were obtained after a subsequent subculturing process. Culture characteristics (on PDA) were observed after three weeks. Type specimens were deposited in the Herbarium of Fungi, Shenzhen University, Shenzhen, China. Living cultures were deposited in the Culture Collection of Microbial Shenzhen University (MBSZU), Shenzhen University, Shenzhen, China. According to Jayasiri et al. (2015) and Index Fungorum (2025), the Faces of Fungi (FOF) and Index Fungorum (IF) numbers were obtained.

DNA extraction, PCR amplification, and sequencing

Fungal cultures grown on PDA for 4 weeks at 25 °C were used for DNA extraction. Axenic mycelium (50–100 mg) was scraped from the growing cultures. The extraction process was performed using a DNA extraction kit (E.Z.N.A. Fungal DNA Mini Kit, D3390-02, Omega Bio-Tek) according to the manufacturer's protocol. In the case of *Acrocalymma ampeli*, *A. poaceicola*, and *Neokalmusia arundinis* (their spores were not germinated), DNA was isolated directly from fungal fruiting bod-

ies on their natural substrate using a commercial DNA extraction kit (E.Z.N.A.® Forensic DNA Kit, D3591-01, Omega BIO-TEK). The DNA products were kept at 4 °C, and replicate samples were stored at -20 °C to ensure long-term stability. The following genomic regions were amplified: the internal transcribed spacer (ITS1–5.8S–ITS2), 28S large subunit rDNA (LSU), 18S small subunit rDNA (SSU), translation elongation factor 1-alpha gene (*tef1-α*), and beta-tubulin (*tub2*). The ITS region was amplified using primers ITS4 and ITS5 (White et al. 1990), the LSU region with LR0R and LR5 (Vilgalys and Hester 1990), and the SSU region with NS1 and NS4 (White et al. 1990). The *tef1-α* gene was amplified using EF1-983F and EF1-2218R (Rehner 2001), and *tub2* was amplified with Bt2a/Bt2b (Glass and Donaldson 1995). The total volume (25 µl) of the PCR reaction contained 9.5 µl of sterilized distilled water, 12.5 µl of 2 × Power Taq PCR MasterMix (a premix and ready-to-use solution, including 0.1 Units/µl Taq DNA Polymerase and 500 µM dNTP Mixture each (dATP, dCTP, dGTP, dTTP) (Bioteke Co., China), 1 µl of each forward and reverse primer (stock concentration 10 pM), and 1 µl of DNA template. Thermal cycle programs for LSU, SSU, ITS, *tef1-α*, and *tub2* genes were followed as mentioned in Tennakoon et al. (2021) and Rathnayaka et al. (2023). The PCR products were quantified using a NanoDrop One spectrophotometer (Thermo Fisher Scientific, USA), verified by 1% agarose gel electrophoresis, and subsequently sequenced by Sangon Biotech (Shanghai) using the original PCR primers.

Phylogenetic analyses

The obtained sequences (forward and reverse) were initially checked with BioEdit v 7.0.5.3 (Hall 1999) and assembled using SeqMan v. 7.0.0 (DNASTAR, Madison, WI). A nucleotide BLAST search (<https://blast.ncbi.nlm.nih.gov/>) was conducted to identify strains with high similarity. Additional related sequences were obtained from recently published data (Tennakoon et al. 2022; Calabon et al. 2023; Konta et al. 2023; Htet et al. 2024). All single-gene sequence datasets were aligned using MAFFT version 7 (Katoh and Standley 2013) through its online platform (<https://mafft.cbrc.jp/alignment/software/>). Single-gene alignment matrices were trimmed to remove ambiguous bases using TrimAl 1.2 (Capella-Gutiérrez et al. 2009). The aligned sequences were concatenated using BioEdit version 7.2.5 (Hall 1999).

The concatenated alignment was subjected to phylogenetic analysis using both Maximum Likelihood (ML) and Bayesian Inference (BI) approaches. Substitution models were selected using MrModeltest v. 2.3 (Nylander 2004) based on the Akaike information criterion (AIC). Maximum Likelihood analysis was performed using RAxML-HPC2 v8.2.8 (Stamatakis 2014) on the CIPRES Science Gateway (Miller et al. 2010), implementing the GTR+I+G model of nucleotide substitution. Bayesian Inference was performed using MrBayes v3.2.1 (Ronquist et al. 2012) with six simultaneous Markov chains run for 3 million generations, sampling trees every 100th generation. The initial 20% of sampled trees were discarded as burn-in, with the remaining 80% used to calculate posterior probabilities and construct the consensus tree. Phylogenetic trees were visualized and initially edited using FigTree v1.4.0 (Rambaut 2012), with final graphical adjustments and layout preparation performed in Microsoft PowerPoint 2010 (Microsoft Corp.) and Adobe Illustrator CS3 (v15.0.0, Adobe Inc.). The newly generated sequences were deposited in GenBank (accession numbers provided in Tables 1–3).

Table 1. Taxa used in the phylogenetic analyses of Acrocalymmaceae and their corresponding GenBank numbers of partial ITS, LSU, and SSU sequences. Isolates/sequences in bold were isolated/sequenced in the present study. Unavailable sequences are indicated as “NA”.

Taxon	Strain / voucher number	GenBank Accession Numbers		
		SSU	LSU	ITS
<i>Acrocalymma ampeli</i>	MFLUCC 20-0159	MW079341	MW063211	MW063150
<i>A. ampeli</i>	NCYUCC 19-0288	MW079342	MW063212	MW063151
<i>A. ampeli</i>	SZU25-013	PV730035	PV730024	PV730045
<i>A. ampeli</i>	SZU25-014	PV730036	PV730025	PV730046
<i>A. aquaticum</i>	MFLUCC 11-0208	JX276953	NG_042698	NR_121544
<i>A. aquaticum</i>	CC36/MFLUCC 20-0124	NA	MT875393	MT875395
<i>A. arengae</i>	MFLUCC 15-0327A	ON650177	ON650673	ON650154
<i>A. arengae</i>	MFLUCC 15-0327B	ON650178	ON650674	ON650155
<i>A. bilobatum</i>	K.L. Chen L119	NA	NA	KX034339
<i>A. bilobatum*</i>	MFLUCC 20-0125	NA	MT875394	MT875396
<i>A. bipolare</i>	MD1321	NA	NG_075326	NA
<i>A. chuxiongense</i>	IFRDCC3104	NA	ON596248	ON595715
<i>A. cycadis</i>	CBS 137972	NA	NG_057046	NR_137884
<i>A. fici</i>	CBS 317.76	NA	NG_057056	NR_137953
<i>A. fici</i>	MFLUCC 21-0103	NA	MT860429	MT864351
<i>A. guizhouense</i>	CGMCC 3.20853	OM838471	OM838474	OM838410
<i>A. guizhouense</i>	GZUIFR H22.028	OM838472	OM838475	OM838411
<i>A. guizhouense</i>	GZUIFR H22.029	OM838473	OM838476	OM838412
<i>A. hongheense</i>	HKAS 111907	MW424792	MW424777	MW424763
<i>A. hongheense</i>	HKAS 111908	MW424791	MW424776	MW424762
<i>A. hongheense</i>	HKAS 111909	MW424790	MW424775	MW424761
<i>A. magnoliae</i>	MFLUCC 18-0545	OL331094	OK655819	OL413439
<i>A. magnoliae</i>	MFLUCC 21-0204	OL331095	OK655820	OL413440
<i>A. medicaginis</i>	CPC 24340	NA	KP170713	KP170620
<i>A. medicaginis</i>	MFLUCC 17-1423	MT214387	MT214432	MT214338
<i>A. medicaginis</i>	MFLUCC 17-1439	MT214388	MT214433	MT214339
<i>A. pterocarpi</i>	MFLUCC 17-0926	MK347840	NG_066306	MK347732
<i>A. pterocarpi</i>	MFLUCC 18-0718	OL331093	OK655818	OL413438
<i>A. pterocarpi</i>	NC13-171	NA	LC517881	LC517880
<i>A. paeoniae</i>	CGMCC:3.24440	OR253217	OR253308	OR253149
<i>A. poaceicola</i>	SZU25-015	PV730032	PV730021	PV730042
<i>A. poaceicola</i>	SZU25-016	PV730033	PV730022	PV730043
<i>A. poaceicola</i>	SZU25-017	PV730034	PV730023	PV730044
<i>A. vagum</i>	CPC 24225	NA	NA	KP170635
<i>A. vagum</i>	CPC 24226	NA	NA	KP170636
<i>A. walkeri</i>	UTHSC DI16-195	NA	LN907338	LT796832
<i>A. yuxiense</i>	HKAS 111910	MW424793	MW424778	NA
<i>Ascocylindrica marina</i>	MD6011	KT252907	KT252905	NA
<i>Ascocylindrica marina</i>	MF416	MK007124	MK007123	NA
<i>Boeremia exigua</i>	CBS 431.74	EU754084	EU754183	FJ427001
<i>Boeremia foveata</i>	CBS 341.67	GU238203	GU237947	GU237834

Table 2. Taxa used in the phylogenetic analyses of Didymosphaeriaceae and their corresponding GenBank numbers of partial ITS, LSU, SSU, and *tef1-α* sequences. Isolates/sequences in bold were isolated/sequenced in the present study. Unavailable sequences are indicated as “NA”.

Taxon	Strain / voucher number	GenBank Accession Numbers			
		LSU	SSU	ITS	<i>tef 1-α</i>
<i>Alloconiothyrium aptrootii</i>	CBS 980.95	JX496234	NA	JX496121	NA
<i>A. aptrootii</i>	CBS 981.95	JX496235	NA	JX496122	NA
<i>A. encephalarti</i>	CPC 35980	MN567610	NA	MN562102	NA
<i>Kalmusia erioi</i>	MFLU 18-0832	MN473052	MN473046	MN473058	MN481599
<i>K. italica</i>	MFLU 14-0620	KP325441	KP325442	KP325440	NA
<i>Laburnicola muriformis</i>	MFLUCC 14-0921	KU743201	KU743202	KU743200	NA
<i>L. muriformis</i>	MFLUCC 16-0290	KU743198	KU743199	KU743197	NA
<i>L. muriformis</i>	MFLUCC 14-0921	KU743201	KU743202	KU743200	NA
<i>L. muriformis</i>	MFLUCC 16-0290	KU743198	KU743199	KU743197	NA
<i>Neokalmusia arundinis</i>	MFLU 17-0754	MT649878	MT649880	MT649882	MT663766
<i>N. arundinis</i>	MFLUCC 15-0463	NG_068237	NG_068372	NR_165852	KY244024
<i>N. arundinis</i>	MFLUCC 14-0222	KX954400	KX986344	KX965731	KY271091
<i>N. arundinis</i>	SZU25-019	PV730028	PV730039	PV730049	PV749905
<i>N. brevispora</i>	KT 2313	AB524601	AB524460	LC014574	AB539113
<i>N. brevispora</i>	KT 1466	AB524600	AB524459	LC014573	AB539112
<i>N. didymospora</i>	MFLUCC 11-0613	KP091434	KP091435	NA	NA
<i>N. jonahhulmei</i>	KUMCC 21-0819	ON007040	ON007049	ON007044	ON009134
<i>N. kunmingensis</i>	KUMCC 18-0120	MK079889	MK079887	MK079886	MK070172
<i>N. scabrispora</i>	KT 1023	AB524593	AB524452	LC014575	AB539106
<i>N. scabrispora</i>	KT 2202	AB524594	AB524453	LC014576	AB539107
<i>N. thailandica</i>	MFLUCC 16-0405	NG_059792	KY706137	NR_154255	KY706145
<i>Neptunomyces aureus</i>	CMG11	NA	NA	MK912120	MK947999
<i>Ne. aureus</i>	CMG12	NA	NA	MK912121	MK948000
<i>Ne. aureus</i>	CMG13	NA	NA	MK912122	MK948001
<i>Ne. aureus</i>	CMG14	NA	NA	MK912123	MK948002
<i>Ne. chinensis</i>	SZU25-020	PV730026	PV730037	PV730047	PV749903
<i>Ne. chinensis</i>	SZU25-021	PV730027	PV730038	PV730048	PV749904
<i>Ne. jeanbriggsiae</i>	BRIP 75897a	PQ431200	NA	PQ431208	NA
<i>Ne. juncicola</i>	CPC 45436	PP791464	NA	PP791436	PP780627
<i>Ne. litoralis</i>	BRIP 75555a	OR259052	NA	OR271911	NA
<i>Ne. soli</i>	MFLUCC 24-0272	PQ306558	NA	NA	NA
<i>Ne. soli</i>	MFLUCC 24-0279	PQ306560	NA	PQ306559	NA
<i>Pseudopithomyces atro-olivaceus</i>	CBS 244.96	LT671616	NA	LT671625	NA
<i>P. chartarum</i>	C459	MK348027	MK347916	MK347808	NA
<i>P. chartarum</i>	C449	MK348017	MK347906	MK347798	NA
<i>P. chartarum</i>	UTHSC 03-2472	HG518064	NA	HG518059	NA
<i>P. chartarum</i>	MFLUCC 17-0314	MF173605	MF173606	MF173607	NA
<i>P. chartarum</i>	C284	MK347964	MK347854	MK347747	NA
<i>P. chartarum</i>	C447	MK348015	MK347904	MK347796	NA
<i>P. chartarum</i>	SZU25-022	PV730030	PV730040	PV730051	PV749906

Taxon	Strain / voucher number	GenBank Accession Numbers			
		LSU	SSU	ITS	<i>tef</i> 1- α
<i>P. diversisporus</i>	UTHSC 06-4528	HG933827	NA	HG933810	NA
<i>P. entadae</i>	C227	MK347946	MK347837	NA	NA
<i>P. karoo</i>	CBS 804.72	NG_057865	HG933811	NR_154291	NA
<i>P. maydicus</i>	MFLUCC 14-0391	KX034666	NA	KX034668	NA
<i>P. maydicus</i>	MFLUCC 17-0028	NA	NA	MG545071	NA
<i>P. palmicola</i>	MFLUCC 14-0392	KX034665	NA	NR_154345	NA
<i>P. sacchari</i>	CBS 803.72	MH872301	LK936379	NA	NA
<i>Spegazzinia</i> sp.	yone 279	AB807583	AB797293	NA	AB808559
<i>Spegazzinia</i> sp.	CL115	AY234948	NA	NA	NA
<i>S. bromeliacearum</i>	URM 8084	MK809513	NA	MK804501	
<i>S. cameliae</i>	CMU 328	MH734521	MH734523	MH734522	MH734524
<i>S. deightonii</i>	yone 212	AB807582	AB797292	NA	AB808558
<i>S. deightonii</i>	MFLUCC 20-0002	MN956772	MN956770	MN956768	MN927133
<i>S. deightonii</i>	yone 66	AB807581	AB797291	NA	AB808557
<i>S. deightonii</i>	MFLUCC 18-1625	ON117309	ON117273	ON117291	ON158097
<i>S. deightonii</i>	SZU25-023	PV730031	PV730041	PV730052	PV749907
<i>S. intermedia</i>	CBS 249.89	MH873861	NA	MH862171	NA
<i>S. lobulata</i>	CBS 361.58	MH869344	NA	MH857812	NA
<i>S. musae</i>	MFLUCC 20-0001	MN930514	MN930513	MN930512	MN927132
<i>S. neosundara</i>	MFLUCC 15-0456	KX954397	KX986341	KX965728	NA
<i>S. neosundara</i>	MFLUCC 13-0211	MH040812	MH040811	MH040810	MH055460
<i>S. tessartha</i>	SH 287	AB807584	AB797294	NA	AB808560
<i>S. tessartha</i>	NRRL 54913	NA	NA	JQ673429	NA
<i>S. tessartha</i>	ASV319	NA	NA	MN898233	NA
<i>S. tessartha</i>	MFLUCC 17-2249	MH071197	MH071192	MH071193	NA
<i>S. tessartha</i>	12H0104	NA	NA	KT385776	NA
<i>S. tessartha</i>	MFLUCC 18-1624	ON117308	ON117272	ON117290	ON158096
<i>S. radermacherae</i>	C264	MK347957	MK347848	MK347740	NA
<i>Xenocamarosporium acaciae</i>	ZHKUCC:22-0202	OP297775	OP297789	OP297805	OP321574
<i>X. acaciae</i>	ZHKUCC:22-0203	OP297776	OP297790	OP297806	OP321575

Results

Phylogenetic analyses

Analysis 1 – Acrocalymmaceae

The combined LSU-SSU-ITS dataset contained 2818 characters, including gaps. *Boeremia exigua* (CBS 431.74) and *B. foveata* (CBS 341.67) were used as outgroup taxa. The RAxML analysis of the combined dataset yielded a best-scoring tree (Fig. 1). The final ML optimization likelihood value was -7357.640966. There were 29.71% undetermined characters or gaps and 509 distinct alignment patterns. Estimated base frequencies were A = 0.250436, C = 0.216202, G = 0.272118, T = 0.261245; substitution rates AC = 1.640471, AG = 1.667992, AT = 2.184081, CG = 0.699183, CT = 5.878108, GT = 1.000; proportion of invariable sites I = 0.692125;

Table 3. Taxa used in the phylogenetic analyses of *Paraconiothyrium* and their corresponding GenBank numbers of partial ITS, LSU, and *tub2* sequences. Isolates/sequences in bold were isolated/sequenced in the present study. Unavailable sequences are indicated as “NA”.

Species	Strain/Voucher No.	GenBank Accession No.		
		LSU	ITS	<i>tub2</i>
<i>Paraconiothyrium ajrekarii</i>	NFCCI 4810	MT372905	MT372906	MT394161
<i>P. archidendri</i>	CBS 168.77	MH872813	MH861045	JX496388
<i>P. archidendri</i>	964-SAB SA1 3	NA	MT820342	NA
<i>P. archidendri</i>	C321	MK347974	MK347757	NA
<i>P. archidendri</i>	1-3-10-2-1-4	NA	KX065269	NA
<i>P. archidendri</i>	NNIBRFG116	NA	KY327413	NA
<i>P. archidendri</i>	NNIBRFG99	NA	KY327412	NA
<i>P. archidendri</i>	NNIBRFG29	NA	KY327411	NA
<i>P. archidendri</i>	MFLUCC 19-0043	ON117302	ON117284	NA
<i>P. archidendri</i>	SZU25-018	PV730029	PV730050	PV749908
<i>P. babiogorense</i>	CBS 128292	MH876291	MH864845	NA
<i>P. brasiliense</i>	CBS 115.92	JX496135	JX496022	JX496361
<i>P. brasiliense</i>	CBS 395.87	JX496196	JX496083	JX496422
<i>P. brasiliense</i>	CBS 122320	JX496146	JX496033	JX496372
<i>P. brasiliense</i>	CBS 122851	JX496149	JX496036	JX496375
<i>P. brasiliense</i>	MFLUCC 19-0040	ON117301	ON117283	NA
<i>P. camelliae</i>	NTUCC 18-096	MT071269	MT112293	MT308623
<i>P. cyclothyrioides</i>	CBS 972.95	JX496232	JX496119	JX496458
<i>P. cyclothyrioides</i>	CBS 432.75	JX496201	JX496088	JX496427
<i>P. cyclothyrioides</i>	SN 3169-19	NA	MN416682	NA
<i>P. cyclothyrioides</i>	NNIBRFG3266	MW237682	NA	NA
<i>P. cyclothyrioides</i>	NNIBRFG3255	MW237679	NA	NA
<i>P. cyclothyrioides</i>	EXF-14614	NA	MT280696	NA
<i>P. cyclothyrioides</i>	NFCCI 4387	MN241143	MN242780	NA
<i>P. estuarinum</i>	CBS 109850	JX496129	JX496016	JX496355
<i>P. fuckelii</i>	JZB320001	MN519513	MN495986	MN508193
<i>P. fuckelii</i>	MFLUCC 13-0073	KJ939281	KJ939278	NA
<i>P. fuckelii</i>	JZB320002	MN519514	MN495987	MN508194
<i>P. fuckelii</i>	JZB320003	MN519515	MN495988	MN508195
<i>P. fuckelii</i>	JZB320004	MN519516	MN495989	MN508196
<i>P. fuckelii</i>	CBS 584.69	NA	JX496211	JX496437
<i>P. fuckelii</i>	CVG970	NA	MZ712975	NA
<i>P. fuckelii</i>	CBS 508.94	JX496209	JX496096	JX496435
<i>P. fuckelii</i>	CBS 653.85	JX496217	JX496104	JX496443
<i>P. fuckelii</i>	CBS 764.71B	JX496225	JX496112	JX496451
<i>P. fuckelii</i>	MFLUCC 19-0067	ON117305	ON117287	NA
<i>P. fuckelii</i>	CBS 797.95	JX496226	JX496113	JX496452
<i>P. fuscomaculans</i>	CBS 116.16	MH866170	MH854649	–

Species	Strain/Voucher No.	GenBank Accession No.		
		LSU	ITS	tub2
<i>P. hakeae</i>	CBS 142521	KY979809	KY979754	KY979920
<i>P. iridis</i>	CBS:146036	MT223919	MT223827	MT223743
<i>P. lini</i>	CBS 253.92	GU238093	NA	KT266268
<i>P. lycopodium</i>	CBS 134705	MH877564	NA	NA
<i>P. maculicutis</i>	CBS 101461	EU754200	NA	NA
<i>P. magnoliae</i>	MFLUCC 10-0278	KJ939283	KJ939280	NA
<i>P. nelloi</i>	MFLU 14-0813	KP711365	KP711360	NA
<i>P. polonense</i>	CBS 134153	KF700360	NA	NA
<i>P. rosae</i>	MFLU 15-1115	MG829041	MG828932	NA
<i>P. thysanolaenae</i>	MFLUCC 10-0550	KP744496	KP744453	NA
<i>P. tiliae</i>	CBS 265.94	EU754139	NA	NA
<i>P. zingiberacearum</i>	MFLUCC 18-0559	ON117303	ON117285	ON158098
<i>P. zingiberacearum</i>	NCYUCC 19-0230	ON117304	ON117286	ON158099
<i>Tremateia arundicola</i>	MFLU 16-1275	KX274248	KX274241	NA
<i>Tremateia guiyangensis</i>	GZAAS01	KX274247	KX274240	NA

gamma distribution shape parameter $\alpha = 0.639926$. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch, respectively (Fig. 1). All analyses (ML and BYPP) yielded similar topologies and concurred with previous studies (Calabon et al. 2023; Konta et al. 2023).

According to the phylogeny, our collection (SZU25-015, SZU25-016, and SZU25-017) clusters within the *Acrocalymma* species. New isolates cluster together and show an independent lineage sister to *A. bilobatum* isolates (MFLUCC 20-0125 and KL Chen L119) with robust statistical support (87% ML, 0.95 BYPP). In addition, the isolates SZU25-013 and SZU25-014 cluster with *A. ampeli* isoates (MFLU 19-2734 and NCYU 19-0008) in a monophyletic clade (100% ML, 1.00 BYPP).

Analysis 2 – *Neokalmusia* and *Neptunomyces*

The combined LSU-SSU-ITS-*tef1*- α dataset contained 3649 characters, including gaps. *Laburnicola muriformis* (MFLUCC 14-0921 and MFLUCC 16-0290) was used as the outgroup taxon. The RAxML analysis of the combined dataset yielded a best-scoring tree (Fig. 2). The final ML optimization likelihood value was -11623.840913. There were 32.16% undetermined characters or gaps and 800 distinct alignment patterns. Estimated base frequencies were A = 0.233832, C = 0.260249, G = 0.273224, T = 0.232696; substitution rates AC = 1.154239, AG = 1.660022, AT = 1.406193, CG = 1.070701, CT = 6.334516, GT = 1.000; proportion of invariable sites I = 0.591573; gamma distribution shape parameter $\alpha = 0.570976$. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch, respectively (Fig. 2). All analyses (ML and BYPP) yielded similar topologies and concurred with previous studies (Crous et al. 2024; Yasanthika et al. 2024).




Figure 1. Phylogram generated from Maximum Likelihood analysis is based on combined LSU, SSU, and ITS sequence data. The tree is rooted with *Boeremia exigua* (CBS 431.74) and *B. foveata* (CBS 341.67). The new isolates are in red, and ex-type strains are indicated in bold face. Bootstrap support values $\geq 65\%$ from the Maximum Likelihood (ML) and Bayesian posterior probabilities (BYPP) values ≥ 0.90 are indicated above the nodes, respectively.

Phylogeny indicates that our collection (SZU25-020 and SZU25-021) clusters within *Neptunomyces* species. Both isolates cluster together and show an independent lineage sister to *N. juncicola* (CPC 45436) with robust statistical support (97% ML, 1.00 BYPP). In addition, the isolate SZU25-019 clusters with *Neokalmusia arundinis* isolates (MFLUCC 15-0463, MFLU 17-0754, and MFLUCC 14-0222) in a monophyletic clade (75% ML, 1.00 BYPP).




Figure 2. Phylogram generated from the Maximum Likelihood analysis is based on combined LSU, SSU, ITS, and *tef1*- α sequence data. The tree is rooted with *Laburnicola muriformis* (MFLUCC 14-0921 and MFLUCC 16-0290). The new isolates are in red, and ex-type strains are indicated in bold face. Bootstrap support values $\geq 65\%$ from the Maximum Likelihood (ML) and Bayesian posterior probabilities (BYPP) values ≥ 0.90 are indicated above the nodes, respectively.

Analysis 3 – *Paraconiothyrium*

The combined LSU-ITS-*tub2* dataset contained 2145 characters, including gaps. *Tremateia arundicola* (MFLUCC 16-1275) and *T. guiyangensis* (GZAAS01) were used as outgroup taxa. The RAxML analysis of the combined dataset yielded the best-scoring tree (Fig. 3). The final ML optimization likelihood value was -7286.035715. There were 35.11% undetermined characters or gaps and 649 distinct alignment patterns. Estimated base frequencies were A = 0.230153, C = 0.249594, G = 0.282309, T = 0.237944; substitution rates AC = 1.589743, AG = 2.770684, AT = 1.318375, CG = 0.871420, CT = 6.249278, GT = 1.000; proportion of invariable sites I = 0.566479; gamma distribution shape




Figure 3. Phylogram generated from the Maximum Likelihood analysis is based on combined LSU, ITS, and *tub2* sequence data. The tree is rooted with *Tremateia arundicola* (MFLUCC 16-1275) and *T. guiyangensis* (GZAAS01). The new isolates are in red, and ex-type strains are indicated in bold face. Bootstrap support values $\geq 65\%$ from the Maximum Likelihood (ML) and Bayesian posterior probabilities (BYPP) values ≥ 0.90 are indicated above the nodes, respectively.

parameter $\alpha = 0.550172$. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch, respectively (Fig. 3). All analyses (ML and BYPP) yielded similar topologies and concurred with previous studies (Guarnaccia et al. 2022; Tennakoon et al. 2022). Phylogeny indicates that our collection (SZU25-018) clusters within *Paraconiothyrium archidendri* isolates (MFLUCC 19-0043, 964-SAB-SA1-3, CBS 168.77, and C321) in a well-supported clade (86% ML, 1.00 BYPP).

Analysis 4 – *Pseudopithomyces*

The combined LSU-SSU-ITS-*tef1*- α dataset comprised 3,215 characters, including gaps. *Spegazzinia radermacherae* (MFLUCC 17-2265) and *S. tessarthra* (SH 287) were used as outgroup taxa. The RAxML analysis of the combined dataset yielded the best-scoring tree (Fig. 4). The final ML optimization likelihood value was -6160.156800. There were 41.67% undetermined characters or gaps and 243 distinct alignment patterns. Estimated base frequencies were A = 0.237009, C = 0.251568, G = 0.272682, T = 0.238741; substitution rates AC = 1.586697, AG = 1.677838, AT = 1.298195, CG = 1.029806, CT = 5.817290, GT = 1.000; proportion of invariable sites I = 0.572938; gamma distribution shape parameter α = 0.783990. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch, respectively (Fig. 4). All analyses (ML and BYPP) yielded similar topologies and concurred with previous studies (Samaradiwakara et al. 2022; Wu et al. 2023). Phylogeny indicates that our collection (SZU25-022) clusters within *Pseudopithomyces chartarum* isolates (C284, C447, C449, C459, UTHSC 04-678, UTHSC 03-2472, and MFLUCC 17-0314) in a well-supported clade (90% ML, 0.98 BYPP).




Figure 4. Phylogram generated from the Maximum Likelihood analysis is based on combined LSU, SSU, ITS, and *tef1*- α sequence data. The tree is rooted with *Spegazzinia radermacherae* (MFLUCC 17-2265) and *S. tessarthra* (SH 287). The new isolates are in red, and ex-type strains are indicated in bold face. Bootstrap support values $\geq 65\%$ from the Maximum Likelihood (ML) and Bayesian posterior probabilities (BYPP) values ≥ 0.90 are indicated above the nodes, respectively.

Analysis 5 – *Spegazzinia*

The combined LSU-SSU-ITS-*tef1-α* dataset contained 3495 characters, including gaps. *Laburnicola muriformis* (MFLUCC 14-0921 and MFLUCC 16-0290) were used as outgroup taxa. The RAxML analysis of the combined dataset yielded the best-scoring tree (Fig. 5). The final ML optimization likelihood value was -8278.609544. There were 35.24% undetermined characters or gaps and 553 distinct alignment patterns. Estimated base frequencies were A = 0.238139, C = 0.257164, G = 0.274729, T = 0.229967; substitution rates AC = 1.482308, AG = 2.141805, AT = 1.664137, CG = 1.084752, CT = 6.606857,




Figure 5. Phylogram generated from the Maximum Likelihood analysis is based on combined LSU, SSU, ITS, and *tef1-α* sequence data. The tree is rooted with *Laburnicola muriformis* (MFLUCC 14-0921 and MFLUCC 16-0290). The new isolates are in red, and ex-type strains are indicated in bold face. Bootstrap support values $\geq 65\%$ from the Maximum Likelihood (ML) and Bayesian posterior probabilities (BYPP) values ≥ 0.90 are indicated above the nodes, respectively.

GT = 1.000; proportion of invariable sites I = 0.670885; gamma distribution shape parameter α = 0.650417. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch, respectively (Fig. 5). All analyses (ML and BYPP) yielded similar topologies and concurred with previous studies (Samarakoon et al. 2020; Tennakoon et al. 2022). Phylogeny indicates that our collection SZU25-023 clusters within *Spegazzinia deightonii* isolates (MFLUCC 20-0002, MFLUCC 18-1625, yone 66, and yone 212) in a well-supported clade (92% ML, 0.95 BYPP).

Taxonomy

Acrocalymma Alcorn & J.A.G. Irwin, Trans. Br. mycol. Soc. 88(2): 163 (1987)

Notes. Alcorn and Irwin (1987) introduced *Acrocalymma* to include *A. medicaginis*, which was recorded as a root pathogen on Medicago in Australia. *Acrocalymma* species show cosmopolitan distribution worldwide (e.g., Australia, China, Egypt, India, Spain, Thailand, and the United States). Their host specificity is also not yet determined and has been recorded from various host families (e.g., Amaranthaceae, Arecaceae, Cucurbitaceae, Cycadaceae, Fagaceae, Lamiaceae, Magnoliaceae, and Moraceae) (Trakunyingcharoen et al. 2014; Jayasiri et al. 2019; Dong et al. 2020; Mortimer et al. 2021; Tennakoon et al. 2021; Calabon et al. 2023; Konta et al. 2023). Most *Acrocalymma* species have been recorded as coelomycetous, and two species have sexual morph, *A. pterocarp*i and *A. hongheense* (Jayasiri et al. 2019; Mortimer et al. 2021). The asexual morph exhibits papillate pycnidia and aseptate, hyaline conidia, which have appendages (Trakunyingcharoen et al. 2014; Jayasiri et al. 2019; Dong et al. 2020), whereas the sexual morph has immersed or semi-immersed, globose to subglobose, ostiolate ascomata, cylindric-clavate asci, and hyaline, fusiform, 1-septate ascospores with distinct sheath (Jayasiri et al. 2019; Mortimer et al. 2021). To date, there are 19 *Acrocalymma* species in Species Fungorum (2025).

Acrocalymma ampeli Tennakoon, C.H. Kuo & K.D. Hyde, Fungal Diversity: 17 (2021)

Index Fungorum: IF555312

Facesoffungi Number: FoF09315

Fig. 6

Description. **Saprobic** on a dead leaf of *Livistona chinensis* (Arecaceae). **Sexual morph:** Undetermined. **Asexual morph:** **Conidiomata** 90–130 × 130–170 µm ($\bar{x}$ = 117 × 153 µm, n = 10), pycnidial, dark brown to black, solitary or clustered, immersed to semi-immersed, erumpent through host surface, unilocular, globose to subglobose, ostiolate. **Conidiomatal wall** 17–23 µm wide ($\bar{x}$ = 19 µm), composed of 4–5 layers of cells with *textura angularis*, cells towards the inside hyaline, and at the outside light brown. **Conidiophores** reduced to conidiogenous cells. **Conidiogenous cells** 5–8 × 4–7 µm ($\bar{x}$ = 6.7 × 5.8 µm, n = 20), hyaline, ampulliform to doliiform, phialidic, smooth-walled. **Conidia** 16–20 × 5–6 µm ($\bar{x}$ = 17.5 × 5.5 µm, n = 20), hyaline, cylindrical to fusoid, apex obtuse, protuber-




Figure 6. *Acrocalymma ampeli* (SZU25-013, new host record). **a, b.** Appearance of conidiomata on the host; **c.** Vertical section of a conidioma; **d, e.** Conidiomatal wall; **f.** Conidiogenous cells with bearing conidia; **g–j.** Conidia; **k, l.** Protruding hila at base (shows in yellow arrows); **m, n.** Mucoid apical appendages (shows in yellow arrows). Scale bars: 50 μm (**c**); 15 μm (**d, e**); 6 μm (**f–j**); 5 μm (**k–n**).

ant and with a rounded hilum at base, aseptate, straight, thin-walled, with flaring mucoid apical appendage at lower end (3–4 μm diam.), visible in water mounts.

Material examined. CHINA • Yunnan Province, Kunming, on a dead leaf of *Livistona chinensis* (Arecaceae), 15 June 2017, D. S. Tennakoon, DST016 (SZU25-013, new host record) • *ibid.* 17 August 2017, DST011 (SZU25-014).

Known hosts. *Livistona chinensis* and *Ficus ampelas* (Tennakoon et al. 2021; this study).

Known distribution. China (Tennakoon et al. 2021; this study).

Notes. *Acrocalymma ampeli* was introduced by Tennakoon et al. (2021) from dead leaves of *Ficus ampelas* (Moraceae). The morphology of our collection (SZU25-013 and SZU25-014) resembles *A. ampeli* by having pycnidial, dark brown to black, immersed to semi-immersed conidiomata, ampulliform to dolii-form conidiogenous cells, and aseptate, cylindrical to fusoid-shaped conidia (Tennakoon et al. 2021). Multi-gene phylogeny also indicates that our collection clusters with *A. ampeli* isolates (MFLUCC 20-0159 and NCYUCC 19-0288) in a well-supported clade (100% ML, 1.00 BYPP). Therefore, based on morphological similarities and phylogeny support, we identified our collection as *A. ampeli* from a different host (*Livistona chinensis*) in China.

***Acrocalymma poaceicola* Tennakoon & S. Hongsanan, sp. nov.**

Index Fungorum: IF903948

Facesoffungi Number: FoF17789

Fig. 7

Etymology. Named after the host family (Poaceae) where this fungus was collected.

Holotype. SZU25-015.

Description. *Saprobic* on a dead leaf of *Arundo pliniana* (Poaceae). **Sexual morph:** Undetermined. **Asexual morph: Conidiomata** 60–70 × 80–110 µm ($\bar{x}$ = 64 × 97 µm, n = 10), pycnidial, dark brown to black, solitary or clustered, immersed to semi-immersed, erumpent through host surface, unilocular, globose to subglobose, ostiolate. **Conidiomatal wall** 15–20 µm wide ($\bar{x}$ = 17 µm), 3–4 layers of irregular cells arranged in a *textura angularis*, cells towards the inside hyaline, at the outside, light brown, thick-walled. **Conidiophores** reduced to conidiogenous cells. **Conidiogenous cells** 5–10 × 2.5–5 µm ($\bar{x}$ = 7.3 × 3.2 µm, n = 20), phialidic, hyaline, smooth,




Figure 7. *Acrocalymma poaceicola* (SZU25-015, holotype). a, b. Appearance of conidiomata on the host; c. Vertical section of a conidioma; d. Conidiomatal wall; e. Conidiogenous cells with developing conidia; f–i. Conidia; j, k. Mucoid apical appendages (shows in yellow arrows); l, m. Protruding hila at base (shows in yellow arrows). Scale bars 30 µm (c); 15 µm (d); 15 µm (e–i); 10 µm (j–m).

ampulliform to doliiform, proliferating with visible periclinal thickening at apex. **Conidia** $24\text{--}32 \times 6\text{--}7.2 \mu\text{m}$ ($\bar{x} = 30 \times 6.8 \mu\text{m}$, $n = 20$), hyaline, cylindrical with an obtuse apex, protuberant and with a rounded hilum at base, straight, aseptate, guttulate, smooth-walled, bearing a mucilaginous appendage ($2\text{--}2.5 \mu\text{m}$ diam.) at the apex.

Material examined. CHINA • Yunnan Province, Kunming, on a dead leaf of *Arundo pliniana* (Poaceae), 12 July 2017, D. S. Tennakoon, DST010 (SZU25-015, **holotype**) • *ibid.* 21 July 2017, DST015 (SZU25-016, **paratype**) • *ibid.* 15 August 2017, DST017 (SZU25-017, **paratype**).

Notes. The morphology of our collection (SZU25-015, SZU25-016, and SZU25-017) tallies with *Acrocalymma* species in having pycnidial conidiomata, hyaline, ampulliform to doliiform conidiogenous cells, and aseptate, cylindrical to fusoid conidia (Zhang et al. 2012; Hyde et al. 2013; Mortimer et al. 2021; Tennakoon et al. 2021). According to the multi-gene phylogeny (LSU, SSU, and ITS) here, our collection clusters with *A. bilobatum* isolates (MFLUCC 20-0125 and K.L. Chen L119) with 87% ML and 0.95 BYPP statistical support. In addition, our collection isolates group together with 100% ML and 1.00 BYPP statistical support. Our collection can be distinguished from *A. bilobatum* in having smaller conidiomata ($60\text{--}70 \times 80\text{--}110 \mu\text{m}$ vs $170\text{--}275 \times 135\text{--}205 \mu\text{m}$) and larger conidia ($24\text{--}32 \times 6\text{--}7.2 \mu\text{m}$ vs $7\text{--}12 \times 2.5\text{--}4 \mu\text{m}$) with bearing a mucilaginous appendage (Calabon et al. 2023). A comparison of the 524 nucleotides across the ITS (+5.8S) gene region of our collection (SZU25-015) and *A. bilobatum* (MFLUCC 20-0125) shows 12 base pair differences (2.29%). Therefore, we recognize that these three isolates belong to one species, which we introduce as a new species herein.

***Neokalmusia* H.A. Ariy. & K.D. Hyde, Fungal Diversity 68: 92 (2014)**

Notes. *Neokalmusia* was introduced by Ariyawansa et al. (2014b) to accommodate two species, *N. brevispora* and *N. scabrispora* (type species), which had been previously classified under *Kalmusia* (Tanaka et al. 2005; Zhang et al. 2009). *Neokalmusia* species have ascomata that are immersed to semi-immersed, globose to subglobose or oblong, with a clypeus-like structure; the asci are cylindric-clavate, with a long pedicel, and the ascospores are fusiform, yellowish-brown to reddish-brown and verrucose (Ariyawansa et al. 2014b; Thambugala et al. 2017; Hongsanan et al. 2020). Currently, ten *Neokalmusia* species are listed in Species Fungorum (2025). Interestingly, all *Neokalmusia* species have a host preference on Poaceae hosts (Ariyawansa et al. 2014b; Thambugala et al. 2017; Hongsanan et al. 2020; Hyde et al. 2020; Wanasinghe et al. 2022). In this study, we introduce a new host record of *N. arundinis* from another Poaceae host, *Sporobolus alterniflorus*.

***Neokalmusia arundinis* Thambugala & K.D. Hyde, Mycosphere 8: 722 (2017)**

Index Fungorum: IF553161

Facesoffungi Number: FoF03219

Fig. 8

Description. **Saprobic** on a dead stem of *Sporobolus alterniflorus* (Poaceae).

Sexual morph: **Ascomata** $300\text{--}370 \times 200\text{--}250 \mu\text{m}$ ($\bar{x} = 340 \times 230 \mu\text{m}$, $n = 6$), immersed, appear as black dots, solitary, shiny, dark brown to black, sub-globose,




Figure 8. *Neokalmusia arundinis* (SZU25-019, new host record). **a** Appearance of ascomata on the host surface; **b**. Close up of an ascoma; **c**. Vertical section of an ascoma; **d**. Section through the peridium; **e**. Pseudoparaphyses; **f–h**. asci; **i–p**. Ascospores. Scale bars: 200 μm (**c**); 15 μm (**d**); 30 μm (**e–h**); 8 μm (**i–p**).

uni-loculate, ostiolate. **Peridium** 14–22 μm wide ($\bar{x}$ = 18 μm), composed of 3–4 layers of brown to dark brown, cells of *textura angularis*, thin-walled. **Hamath-ecium** comprising 1.5–3 μm wide, numerous, cellular, pseudoparaphyses. **Asci** 75–95 $\times$ 7–9 μm ($\bar{x}$ = 85 $\times$ 8 μm , n = 20), 8-spored, bitunicate, fissitunicate, cylindric-clavate, long pedicellate, furcate at base, apically rounded with an indistinct ocular chamber. **Ascospores** 13–17 $\times$ 3.8–5.2 μm ($\bar{x}$ = 15 $\times$ 4.2 μm , n = 30), overlapping, 1–2-seriate, hyaline when immature, pale brown to dark brown at maturity, fusiform, 1-septate, distinctly constricted at the septum, straight or slightly curved, asymmetrical, upper cell shorter than lower cell, often enlarged near septum in the upper cell, smooth-walled, **Asexual morph**: Undetermined.

Material examined. CHINA • Yunnan Province, Kunming, on dead stem of *Sporobolus alterniflorus* (Poaceae), 28 December 2024, D.S. Tennakoon, DSZ21 (SZU25-019, new host record).

Known hosts. *Arundo pliniana*, *Panicum virgatum*, *Sporobolus alterniflorus* (Thambugala et al. 2017; Hongsanan et al. 2020; this study).

Known distribution. China and Italy (Thambugala et al. 2017; Hongsanan et al. 2020; this study).

Notes. *Neokalmusia arundinis* was introduced by Thambugala et al. (2017) from a dead stem of *Arundo pliniana* in Italy. The morphology of our collection (SZU25-019) tallies well with the type of *N. arundinis* (MFLU 16-2577) by having immersed, globose to sub-globose, scattered, dark brown ascomata with a clypeus, cylindric-clavate, long pedicellate asci, and fusiform, pale brown to brown, 1-septate ascospores (Thambugala et al. 2017). In addition, the sizes of ascomata ($275\text{--}350 \times 225\text{--}275 \mu\text{m}$ vs $300\text{--}370 \times 200\text{--}250 \mu\text{m}$), asci ($60\text{--}85 \times 8.5\text{--}10.5 \mu\text{m}$ vs $75\text{--}95 \times 7\text{--}9 \mu\text{m}$), and ascospores ($11.8\text{--}16.2 \times 4\text{--}5.4 \mu\text{m}$ vs $13\text{--}17 \times 3.8\text{--}5.2 \mu\text{m}$) also overlap between our collection and the type of *N. arundinis* (MFLU 16-2577). According to the multi-gene phylogeny, our collection groups with other *N. arundinis* isolates in a well-supported clade (75% ML, 1.00 BYPP). Therefore, we introduce our collection as a new host record of *N. arundinis* from *Sporobolus alterniflorus* in China.

***Neptunomyces* M. Gonçalves, T. Vicente & A. Alves, MycoKeys 60: 37 (2019)**

Notes. Gonçalves et al. (2019) introduced *Neptunomyces*, which includes *N. aureus*, isolated from *Ulva* sp. in Portugal. *Neptunomyces* species have aseptate, golden yellow, subcylindrical conidia with rounded apices. Currently, five *Neptunomyces* species are listed in Index Fungorum (2025): *N. aureus*, *N. jeanbriggsiae*, *N. juncicola*, *N. litoralis*, and *N. soli*. In this study, we introduce another *Neptunomyces* species, *N. chinensis*, from *Phragmites australis* in China.

***Neptunomyces chinensis* Tennakoon & S. Hongsanan, sp. nov.**

Index Fungorum: IF903947

Facesoffungi Number: FoF17788

Fig. 9

Etymology. Named after the country (China) where this fungus was collected.

Holotype. SZU25-020.

Description. **Saprobic** on dead leaf of *Phragmites australis* (Poaceae). **Sexual morph:** Undetermined. **Asexual morph:** Coelomycetous. **Conidiomata** $90\text{--}110 \times 80\text{--}150 \mu\text{m}$ ($\bar{x} = 102 \times 120 \mu\text{m}$, $n = 10$), pycnidial, immersed to semi-immersed, dark brown to black, solitary to aggregated, globose to sub-globose, visible as dots on host surface. **Conidiomatal wall** $15\text{--}20 \mu\text{m}$ wide ($\bar{x} = 16 \mu\text{m}$), thick-walled, composed of several layers of brown to dark brown pseudoparenchymatous cells, fusing at the outside indistinguishable from the host tissues. **Conidiophores** reduced to conidiogenous cells. **Conidiogenous cells** $4\text{--}7 \times 3\text{--}4 \mu\text{m}$ ($\bar{x} = 4.8 \times 3.5 \mu\text{m}$, $n = 20$), discrete, hyaline, globose to doliiform, holoblastic. **Conidia** $6.5\text{--}8 \times 4\text{--}5$ ($\bar{x} = 6.8 \times 4.5 \mu\text{m}$, $n = 40$) μm , hyaline to pale brown, ellipsoidal to limoniform, apex acute to apiculate, widest in the middle, tapering towards a narrowly truncate base, guttulate, smooth-walled.

Material examined. CHINA • Guangdong Province, Shenzhen, on a dead leaf of *Phragmites australis* (Poaceae), 20 December 2024, D.S. Tennakoon, SZD 12 (SZU25-020, **holotype**), ex-type living culture, MBSZU 25-028 • *ibid.* 23 December 2024, SZD15 (SZU25-021, **paratype**), living culture, MBSZU 25-029.




Figure 9. *Neptunomyces chinensis* (SZU25-020, holotype). a, b. Appearance of conidiomata on host; c. Close-up of conidioma; d, e. Vertical sections of conidiomata; f. Conidiomatal wall; g. Conidiogenous cells with developing conidia; h, i. Conidia. Scale bars: 60 μm (d, e); 15 μm (f); 5 μm (g–i).

Notes. According to the multi-gene phylogeny, our collection (SZU25-020 and SZU25-021) constitutes an independent lineage sister to *Neptunomyces juncicola* (CPC 45436) with robust statistical support (97% ML, 1.00 BYPP). In addition, new isolates cluster together with strong support (100% ML, 1.00 BYPP). Our collection can be distinguished from *N. juncicola* by conidial characteristics; *N. juncicola* has subcylindrical to fusoid-ellipsoid conidia with sub-obtuse apex, whereas our collection has ellipsoidal to limoniform conidia with acute to apiculate apex (Crous et al. 2024). Our collection differs from *N. juncicola* by having smaller conidiomata (90–110 $\times$ 80–150 μm vs 180–220 μm). However, SZU25-020 and SZU25-021 have some morphological similarities with *N. soli* (immersed to semi-immersed, globose to sub-globose conidiomata and dolii-form conidiogenous cells and hyaline to pale brown conidia), but differ by conidia sizes (6.5–8 $\times$ 4–5 μm vs 5–6.5 $\times$ 3–5 μm) (Yasanthika et al. 2024). In addition, a comparison of the 587 nucleotides across the ITS (+5.8S) gene region of our collection (SZU25-020) and *N. soli* (MFLUCC 24-0272) shows 19 base pair differences (3.23%). Therefore, based on both morphological and phylogenetic evidence, we introduce our collection as a new species, *N. chinensis*, from China.

***Paraconiothyrium* Verkley, Stud. Mycol. 50(2): 327 (2004)**

Notes. Verkley et al. (2004) introduced *Paraconiothyrium*, which includes four species: *P. estuarinum* (type species), *P. brasiliense*, *P. cyclothyrioides*, and *P. fungicola*. *Paraconiothyrium* species have variable morphological characteristics, including eustromatic to pycnidial conidiomata, phialidic or annelidic conidiogenous cells, smooth-walled or minutely warted, hyaline or brown conidia (Verkley et al. 2004; Damm et al. 2008; Paul and Lee 2014). The life modes of *Paraconiothyrium* can vary, as they are primarily reported as saprobes, with some species also being identified as endophytes and pathogens (Khan et al. 2012; Nakashima et al. 2019; Wang et al. 2021; Guarnaccia et al. 2022; Tennakoon et al. 2022). They also have diverse bioactive functions with potential applications in agriculture, medicine, and industrial sectors (Wang et al. 2021). To date, 28 species of *Paraconiothyrium* have been listed in Species Fungorum (2025). In this study, we introduce a new host record of *P. archidendri* from *Citrus maxima* in China.

***Paraconiothyrium archidendri* Verkley, Göker and Stielow, Persoonia 32: 37 (2014)**

Index Fungorum: IF800761

Facesoffungi Number: FoF05243

Fig. 10

Description. **Saprobic** on a dead stem of *Citrus maxima* (Rutaceae). **Sexual morph:** Undetermined. **Asexual morph:** Coelomycetous. **Conidiomata** 100–150 × 100–170 µm ($\bar{x}$ = 105 × 135 µm, n = 10), pycnidial, immersed to semi-immersed, dark brown to black, solitary to aggregated, globose to sub-globose. **Conidiomatal wall** 15–30 µm wide ($\bar{x}$ = 17 µm), thick-walled, composed of several layers of brown to dark brown pseudoparenchymatous cells, cells towards the inside hyaline, arranged in a *textura angularis*, fusing at the outside indistinguishable from the host tissues. **Conidiophores** reduced to conidiogenous cells. **Conidiogenous cells** 6–8 × 3–4 µm ($\bar{x}$ = 6.4 × 3.2 µm, n = 20), discrete, hyaline, globose to doliiform, holoblastic. **Conidia** 5.8–7 × 3–4 ($\bar{x}$ = 6.3 × 3.2 µm, n = 40) µm, variable in shape, subglobose to ellipsoid, rarely obovoid, ends rounded, aseptate, initially hyaline, becoming olivaceous-brown at maturity, contents with several small oil droplets, smooth-walled.

Known hosts. *Acer pentaphyllum*, *Alliaria petiolate*, *Cinnamomum camphora*, *Citrus maxima*, *Coffea arabica*, *Ginkgo biloba*, *Magnolia* sp., *Picea glauca*, *Pinus tabulaeformis*, *Pithecolobium bigeminum* and *Prunus salicina* (Verkley et al. 2004, 2014; Damm et al. 2008; Paul and Lee 2014; Nakashima et al. 2019; Tennakoon et al. 2022; this study).

Known distribution. Brazil, Canada, China, Japan, Korea, Myanmar, South Africa, and the United States (Verkley et al. 2004, 2014; Damm et al. 2008; Paul and Lee 2014; Nakashima et al. 2019; Tennakoon et al. 2022; this study).

Material examined. CHINA • Guangdong Province, Shenzhen, on dead stem of *Citrus maxima* (Rutaceae), 21 December 2024, D. S. Tennakoon, DC030 (SZU25-018, new host record); living culture MBSZU 25-027.

Notes. *Paraconiothyrium archidendri* was initially introduced by Verkley et al. (2014) from *Pithecellobium bigeminum* in Myanmar. A multi-gene phylogeny in-




Figure 10. *Paraconiothyrium archidendri* (SZU25-018, new host record). a. Conidiomata on a dead stem of *Citrus maxima*; b, c. Close-up of conidiomata on the host; d. Section through conidioma; e. Conidiomatal wall; f. Conidiogenous cells with developing conidia; g–i. Conidia. Scale bars: 50 μ m (d); 20 μ m (e); 5 μ m (f–i).

indicates that our collection (SZU25-018) groups with other *P. archidendri* isolates, particularly closely related to the isolate CBS 168.77. Our collection shares similar morphological characteristics with *P. archidendri* by having pycnidial, immersed to semi-immersed, dark brown to black conidiomata, globose to doliiform conidiogenous cells, and subglobose to ellipsoid, olivaceous-brown, aseptate conidia (Damm et al. 2008; Paul and Lee 2014; Verkley et al. 2014; Nakashima et al. 2019; Tennakoon et al. 2022). Thus, we identify our collection as *Paraconiothyrium archidendri* with a new host occurrence from *Citrus maxima* in China.

***Pseudopithomyces* Ariyaw. & K.D. Hyde, Fungal Diversity 75: 64 (2015)**

Notes. Ariyawansa et al. (2015) introduced this genus to place *Pseudopithomyces chartarum* as the type species. *Pseudopithomyces* has a cosmopolitan distribution worldwide (Hyde et al. 2017; Wu et al. 2023; Farr and Rossman 2025). Species have fusiform, verruculose, dark conidia and produce brown to black colonies on the host (Ariyawansa et al. 2015; Jayasiri et al. 2019; Tenna-

koon et al. 2021). Currently, 13 *Pseudopithomyces* species are listed in Species Fungorum (2025). In this study, we introduce a new host record of *P. chartarum* from a dead leaf of *Hedychium coronarium* (Zingiberaceae).

***Pseudopithomyces chartarum* (Berk. & M.A. Curtis) Jun F. Li, Ariyaw. & K.D. Hyde, Fungal Divers 75: 66 (2015)**

Index Fungorum: IF551393

Facesoffungi Number: FoF00938

Fig. 11

Description. **Saprobic** on a dead leaf of *Hedychium coronarium* (Zingiberaceae). **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. Colonies effuse, scattered, powdery, dark brown to black. **Vegetative hyphae** superficial or partly immersed in the substrate, composed of septate, branched, smooth, thin-walled hyphae. **Conidiophores** mononematous, micronematous, mostly intercalary, sometimes denticulate, aseptate. **Conidiogenous cells** with 2 µm broad conidial attachment, terminal, hyaline, globose or subglobose, integrated, hyaline to pale brown. **Conidia** 15–21 × 8–10 µm ($\bar{x}$ = 17 × 9.1 µm, n = 40), solitary, initially light brown, becoming brown to dark brown at maturity, obovate to oblong, verruculose to spinulose, muriform, 3–4 vertical septa, mostly 1–2 longitudinal dark septa, darken and slightly constricted at the septa, thick-walled.

Material examined. CHINA • Guangdong Province, Shenzhen, on dead leaf of *Hedychium coronarium* (Zingiberaceae), 15 December 2024, D.S. Tennakoon, DSZ10 (SZU25-022, new host record), living culture, MBSZU 25-030.

Known hosts. *Bauhinia* spp., *Centrosema pubescens*, *Ceratonia siliqua*, *Chloris gayana*, *Chrysanthemum coronarium*, *Cirsium arvense*, *Cocos* spp., *Colocasia* spp., *Commelina benghalensis*, *Conyza bonariensis*, *Crotalaria pseudospartium*, *Crotalaria striata*, *Cunninghamia lanceolata*, *Cupania macrophylla*, *Cynodon* spp., *Dactylis glomerata*, *Daucus* spp., *Descurainia sophia*, *Desmodium* spp., *Digitaria* spp., *Dolichos* spp., *Eichhornia crassipes*, *Elegia* spp., *Eucalyptus* spp., *Foeniculum vulgare*, *Glycine* spp., *Gossypium* spp., *Guazuma ulmifolia*, *Harpullia* spp., *Ipomoea* spp., *Juncus roemerianus*, *Leucaena* spp., *Macaranga tanarius*, *Magnolia grandiflora*, *Malus* spp., *Medicago* spp., *Miscanthus* spp., *Morus* spp., *Musa* spp., *Oryza sativa*, *Pandanus tectorius*, *Panicum* spp., *Phyllostachys* spp., *Radermachera sinica*, and *Zea mays* (Hyde et al. 2017; Jayasiri et al. 2019; Tennakoon et al. 2021; Farr and Rossman 2025; this study).

Known distribution. Australia, Austria, Brazil, Canada, China, Cuba, Japan, Ghana, Greece, India, Indonesia, Kenya, Malaysia, Myanmar, Nicaragua, New Zealand, Papua New Guinea, Poland, South Africa, Sudan, Thailand, Venezuela, the United States (Hyde et al. 2017; Jayasiri et al. 2019; Tennakoon et al. 2021; Farr and Rossman 2025; this study).

Notes. Due to the morphological characteristics largely overlapping with those of *Pseudopithomyces chartarum* isolates, we report our collection (SZU25-022) as a new host record of *P. chartarum* from dead leaves of *Hedychium coronarium* (Zingiberaceae). For instance, our collection resembles *P. chartarum*, having powdery, dark brown to black colonies on the host surface and brown, obovate to oblong, muriform conidia (Ariyawansa et al. 2015; Jayasiri et




Figure 11. *Pseudopithomyces chartarum* (SZU25-022, new host record). **a–c.** Appearance of colonies on host surface; **d–g.** Conidiogenous cells bearing conidia; **h–m.** Conidia. Scale bars: 10 µm (d–m).

al. 2019; Samaradiwakara et al. 2022; Wu et al. 2023). Phylogeny also shows that our collection groups with other *P. chartarum* isolates in a well-supported clade (90% ML, 0.98 BYPP).

***Spegazzinia* Sacc. Padova: [1] (1879)**

Notes. *Spegazzinia* is a diverse, hyphomycetous genus introduced by Saccardo (1880) to include *S. ornata* as the type species. *Spegazzinia* species have been reported mainly as saprobes on litter of tropical, subtropical, and temperate vascular plants (Matsushima 1980; Lu et al. 2000; Manoharachary and Kunwar 2010; Thambugala et al. 2017). Additionally, some have been identified as endophytes (e.g., *S. tessarthra* and *S. bromeliacearum*) (Manish et al. 2014; Crous et al. 2019). Apart from that, *Spegazzinia* species have also been found in soil (Ellis 1971). The morphological characteristics of *Spegazzinia* are quite distinct from those of other hyphomycetous genera due to its pleomorphism (Mena-Portales et al. 2017). They have two types of conidia in the same mycelium, such as α and β conidia. Some species have spines in both types of conidia, while some taxa do not bear spines. Currently, 28 *Spegazzinia* species are listed in Species Fungorum (2025). In this study, we introduce a new host record of *S. deightonii* from a dead leaf of *Arundo pliniana* in China.

***Spegazzinia deightonii* (S. Hughes) Subram., J. Indian Bot. Soc. 35: 78 (1956)**

Index Fungorum: IF306062

Facesoffungi Number: FoF07238

Fig. 12

Description. *Saprobic* on a dead leaf of *Arundo pliniana* (Poaceae). **Sexual morph:** Undetermined. **Asexual morph:** Hyphomycetous. **Sporodochia** 1–2 mm diam., dark, black, dense, powdery, velvety. **Conidiophores** give rise to two types of conidia referred to here as α and β . **Conidiophores** of α conidia up to $75\text{--}90 \times 1\text{--}2 \mu\text{m}$ ($\bar{x} = 80 \times 1.6 \mu\text{m}$, $n = 20$) long, erect or flexuous, narrow, verruculose, unbranched, base light brown, upper part dark brown. **Conidiogenous cell** development basauxic, forming a single, terminal holoblastic conidium at the apex of the conidiophore. Conidial development holoblastic. **Conidia** two types: **α conidia** $15\text{--}27 \times 16\text{--}24 \mu\text{m}$ ($\bar{x} = 25 \times 21 \mu\text{m}$, $n = 25$), stellate, solitary, globose to variously shaped, with spines $4\text{--}6 \mu\text{m}$ long, 4–8-celled,




Figure 12. *Spegazzinia deightonii* (SZU25-023, new host record). a. Sporodochia on a dead leaf of *Arundo pliniana*; b, c. Close-up of sporodochia; d–f. α Conidia with conidiophores; g, h. α Conidia with conidiogenous cells; i–m. β Conidia. Scale bars: $15 \mu\text{m}$ (d–h); $10 \mu\text{m}$ (i–m).

deeply constricted at the septa. ***β conidia*** 16–22 × 10–15 µm ($\bar{x}$ = 19 × 14 µm, n = 25), disc-shaped, initially hyaline, light brown to dark brown at maturity, 8-celled, flat from both sides, frequently with attached conidiogenous cells when splitting from the conidiophores.

Known hosts. *Arundo pliniana*, *Areca catechu*, *Cocos nucifera*, *Hedychium coronarium*, *Musa* sp., and *Panicum maximum* (Matsushima 1980; Lu et al. 2000; Tianyu 2009; Samarakoon et al. 2020; Tennakoon et al. 2022; Farr and Rossman 2025; this study).

Known distribution. China and Thailand (Matsushima 1980; Lu et al. 2000; Tianyu 2009; Samarakoon et al. 2020; Tennakoon et al. 2022; Farr and Rossman 2025; this study).

Material examined. CHINA • Guangdong Province, Shenzhen, on dead leaf of *Arundo pliniana* (Poaceae), 21 December 2024, D. S. Tennakoon, DROD009 (SZU25-023, new host record); living culture, MBSZU 25-031.

Notes. The morphological characteristics of our collection (SZU25-023) resemble those of *Spegazzinia deightonii*, as it has 8-celled, disk-shaped, dark brown, spiny conidia (Tanaka et al. 2015; Samarakoon et al. 2020; Tennakoon et al. 2022). Phylogeny also indicates that our collection clusters with other *S. deightonii* isolates in a well-supported clade (92% ML, 0.95 BYPP), and makes a close phylogenetic relationship with the isolate MFLUCC 20-0002 (96% ML, 0.96 BYPP). Therefore, we identify our collection as an isolate of *S. deightonii* from a dead leaf of *Arundo pliniana* in China.

Discussion

Fungal taxonomy has undergone significant evolution over the last two decades, transitioning from traditional morphology-based classification to the current era of DNA-based fungal classification (Tedersoo et al. 2020; Lücking et al. 2021). Initially, morphological characteristics served as the primary criteria for fungal classification, which were mostly represented by line drawings, taxonomic keys, descriptions, and illustrations (Ellis 1971; Nagraj 1978; Sutton 1980). The traditional approach persisted for centuries and was largely limited by factors such as phenotypic plasticity, which has led to misinterpretations and taxonomic confusions (Alster et al. 2021). Consequently, morphological characteristics combined with a DNA-based taxonomic approach have emerged as the main research tool for fungal classification and have facilitated the resolution of species boundaries with numerous novel fungal discoveries (White et al. 1990; Tanaka et al. 2015; Hyde et al. 2020; Crous et al. 2021; Mortimer et al. 2021; Tennakoon et al. 2024). In particular, molecular techniques have facilitated the construction of fungal phylogenies, the identification of evolutionary patterns, and the discovery of cryptic species (Hyde et al. 2017; Fontecha et al. 2019; Hongsanan et al. 2020). However, the accurate identification of fungal species remains a significant challenge. To overcome these challenges, mycologists have made significant strides in resolving fungal taxonomy across various taxonomic levels, including orders, families, genera, and species. For instance, mycologists have identified the class Dothideomycetes, comprising 46 orders, 231 families, 2,346 genera, and 32,621 species (Hongsanan et al. 2020; Wijayawardene et al. 2022; Bánki et al. 2023).

We describe and illustrate two new fungal species (*Acrocalymma poaceicola* and *Neptunomyces chinensis*) and five new host records (*A. ampeli*,

Neokalmusia arundinis, *Paraconiothyrium archidendri*, *Pseudopithomyces chartarum*, and *Spegazzinia deightonii*) in Acrocalymmaceae and Didymosphaeriaceae families (Pleosporales, Dothideomycetes) based on morphological and molecular analyses (Figs 6–12). All species were collected from forest plant litter. The newly proposed species, *Acrocalymma poaceicola* and *Neokalmusia chinensis*, are distinguished from related taxa through both distinct morphological characteristics and molecular phylogenetic evidence. Additionally, the morphological features of the new host records align with those of their respective type specimens, exhibiting dimensional overlap.

The species of Didymosphaeriaceae have a cosmopolitan distribution worldwide and have been recorded from a vast range of host families (Ariyawansa et al. 2014a, b; Hyde et al. 2013; Liu et al. 2015; Wanasinghe et al. 2016; Phookamsak et al. 2019; Tennakoon et al. 2022; Farr and Rossman 2025). Of them, some Didymosphaeriaceae genera are vastly diverse with many species (e.g., *Didymosphaeria*: 195 species, *Paraphaeosphaeria*: 40 species, *Montagnula*: 49 species, *Kalmusia*: 31 species, *Paraconiothyrium*: 28 species, *Karstenula*: 23 species, *Phaeodothis*: 22 species, *Spegazzinia*: 28 species, *Pseudocamarosporium*: 15 species, *Pseudopithomyces*: 15 species, *Letendraea*: 12 species, *Neokalmusia*: 10 species, *Tremateia*: 10 species, *Paracamarosporium*: 9 species, *Chromolaenicola*: 8 species, *Laburnicola*: 8 species, *Deniquelata*: 7 species, *Julella*: 7 species, *Curreya*: 6 species, *Neptunomyces*: 5 species). Some genera consist of less than 5 species (e.g., *Alloconiothyrium*: 3 species, *Austropleospora*: 3 species, *Verrucoconiothyrium*: 3 species, *Bimuria*: 2 species, *Cylindroaseptospora*: 2 species, *Didymocrea*: 2 species, *Paramassariosphaeria*: 2 species). In addition, some genera remain monotypic, thus additional collections are needed to facilitate their expansion (e.g., *Barria*, *Kalmusibambusa*, *Lineostroma*, *Vicosamyces*, *Xenocamarosporium*). Despite having a large number of genera in Didymosphaeriaceae, some are still lacking molecular data (e.g., *Barria*, *Lineostroma*). Thus, future collections are essentially required to resolve their phylogenetic relationships in Didymosphaeriaceae.

By contrast, Acrocalymmaceae is monotypic, consisting only of *Acrocalymma*, which has 19 species. Most of the Acrocalymmaceae taxa play a crucial role in various ecological niches (e.g., terrestrial, freshwater, soil) as saprobes (Trakunyingcharoen et al. 2014; Jayasiri et al. 2019; Mortimer et al. 2021; Tennakoon et al. 2021; Konta et al. 2023). In addition, they are known to act as pathogens as well (e.g., *A. medicaginis*, *A. vagum*) (Trakunyingcharoen et al. 2014). The host specificity of Acrocalymmaceae still needs to be investigated, as they have been reported from a wide host range (e.g., *Anomianthus dulcis*, *Arenga pinnata*, *Cycas calcicole*, *Ficus ampelas*, *Magnolia* spp., and *Quercus glauca*) (Trakunyingcharoen et al. 2014; Jayasiri et al. 2019; Das et al. 2020; Mortimer et al. 2021; Tennakoon et al. 2021; Calabon et al. 2023; Konta et al. 2023). Nonetheless, it is worthwhile to collect additional fungal samples from various habitats and ecological niches to facilitate the continued expansion of this genus.

Acknowledgments

The authors would like to thank Shaun Pennycook (Manaaki Whenua Landcare Research, New Zealand) for his guidance on the fungal nomenclature. Danushka S.

Tennakoon would like to thank the Postdoctoral Fellowship in Shenzhen Key Laboratory of Microbial Genetic Engineering, College of Life Science and Oceanography, Shenzhen University, Shenzhen 518060, China.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

This research was funded by Natural Science Foundation of Guangdong Province (2024B1515020034), and National Key R&D Program of China (2021YFA0910800).

Author contributions

Conceptualization, D.S.T., N.I.d.S. and S.H.; methodology, D.S.T. and N.I.d.S.; software, D.S.T. and N.I.S.; validation, N.I.d.S., S.H. and N.X.; formal analysis, D.S.T. and N.I.d.S.; investigation, D.S.T., N.I.d.S., S.H. and N.X.; resources, D.S.T., S.H. and N.X.; data curation, D.S.T. and N.I.d.S.; writing – original draft preparation, D.S.T., N.I.d.S., S.H. and N.X.; writing – review and editing, D.S.T., N.I.d.S., S.H. and N.X.; supervision, S.H. and N.X.; project administration, S.H. and N.X.; funding acquisition, S.H. and N.X. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Danushka S. Tennakoon  <https://orcid.org/0000-0003-2306-1255>

Nimali I. de Silva  <https://orcid.org/0000-0002-1577-280X>

Sinang Hongsanant  <https://orcid.org/0000-0003-0550-3152>

Ning Xie  <https://orcid.org/0000-0002-5866-8535>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Alcorn JL, Irwin JAG (1987) *Acrocalymma medicaginis* gen. et sp. nov. causing root and crown rot of *Medicago sativa* in Australia. Transactions of the British Mycological Society 88(2): 163–167. [https://doi.org/10.1016/S0007-1536\(87\)80211-5](https://doi.org/10.1016/S0007-1536(87)80211-5)
- Alster CJ, Allison SD, Johnson NG, Glassman SI, Treseder KK (2021) Phenotypic plasticity of fungal traits in response to moisture and temperature. ISME Communications 1(1): 43. <https://doi.org/10.1038/s43705-021-00045-9>
- Ariyawansa HA, Camporesi E, Thambugala KM, Mapook A, Kang JC, Alias SA, Chukeatirote E, Thines M, Mckenzie EH, Hyde KD (2014a) Confusion surrounding Didymosphaeria—Phylogenetic and morphological evidence suggest Didymosphaeriaceae is not a distinct family. Phytotaxa 176(1): 102–119. <https://doi.org/10.11646/phytotaxa.176.1.12>

- Ariyawansa HA, Tanaka K, Thambugala KM, Phookamsak R, Tian Q, Camporesi E, Hongsan S, Monkai J, Wanasinghe DN, Mapook A, Chukeatirote E, Kang JC, Xu JC, McKenzie EHC, Jones EBG, Hyde KD (2014b) A molecular phylogenetic reappraisal of the Didymosphaeriaceae (=Montagnulaceae). *Fungal Diversity* 68: 69–104. <https://doi.org/10.1007/s13225-014-0305-6>
- Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B, Chethana KWT, Dai DQ, Dai YC, Daranagama DA, Jayawardena RS, Lücking R, Ghobad-Nejhad M, Niskanen T, Thambugala KM, Voigt K, Zhao RL, Li GJ, Doilom M, Boonmee S, Yang ZL, Cai Q, Cui YY, Bahkali AH, Chen J, Cui BK, Chen JJ, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, Hashimoto A, Hongsan S, Jones EBG, Larsson E, Li WJ, Li QR, Liu JK, Luo ZL, Maharachchikumbura SSN, Mapook A, McKenzie EHC, Norphanphoun C, Konta S, Pang KL, Perera RH, Phookamsak R, Phukhamsakda C, Pinruan U, Randrianjohany E, Singtripop C, Tanaka K, Tian CM, Tibpromma S, Abdel-Wahab MA, Wanasinghe DN, Wijayawardene NN, Zhang JF, Zhang H, Abdel-Aziz FA, Wedin M, Westberg M, Ammirati JF, Bulgakov TS, Lima DX, Callaghan TM, Callac P, Chang CH, Coca LF, Dal-Forno M, Dollhofer V, Fliegerová K, Greiner K, Griffith GW, Ho HM, Hofstetter V, Jeewon R, Kang JC, Wen TC, Kirk PM, Kytövuori I, Lawrey JD, Xing J, Li H, Liu ZY, Liu XZ, Liimatainen K, Lumbsch HT, Matsumura M, Moncada B, Nuankaew S, Parnmen S, de Azevedo SALCM, Sommai S, Song Y, de Souza CAF, de Souza-Motta CM, Su HY, Suetrong S, Wang Y, Wei SF, Wen TC, Yuan HS, Zhou LW, Réblová M, Fournier J, Camporesi E, Luangsa-ard JJ, Tasanathai K, Khonsanit A, Thanakitpipattana D, Somrithipol S, Diederich P, Millanes AM, Common RS, Stadler M, Yan JY, Li XH, Lee HW, Nguyen TTT, Lee HB, Battistin E, Marsico O, Vizzini A, Vila J, Ercole E, Eberhardt U, Simonini G, Wen HA, Chen XH, Miettinen O, Spirin V, Hernawati (2015) Fungal diversity notes 111–252—Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 75: 27–274. <https://doi.org/10.1007/s13225-015-0346-5>
- Asplund J, Kauserud H, Bokhorst S, Lie MH, Ohlson M, Nybakken L (2018) Fungal communities influence decomposition rates of plant litter from two dominant tree species. *Fungal Ecology* 32: 1–8. <https://doi.org/10.1016/j.funeco.2017.11.003>
- Bánki O, Roskov Y, Döring M, Ower G, Vandepitte L, Hobern D, Remsen D, Schalk P, De Walt RE, Keping M (2023) Catalogue of Life Checklist (Version 2023-01-12). Catalogue of Life. <https://doi.org/10.48580/dfqz>
- Barr ME (1990) Melanommatales (Loculoascomycetes). *North American Flora* 13: 1–129.
- Berg B, McClaugherty C (2008) Plant litter: decomposition, humus formation, carbon sequestration. Springer, Berlin.
- Calabon MS, Jones EG, Boonmee S, Li WJ, Xiao YP, Hyde KD (2023) Acrocalymmaceae (Pleosporales) from freshwater habitats in Thailand with the introduction of *Acrocalymma bilobatum* sp. nov. *Studies in Fungi* 8: 11. <https://doi.org/10.48130/SIF-2023-0011>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) TrimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics (Oxford, England)* 25: 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Crous PW, Carnegie AJ, Wingfield MJ, Sharma R, Mughini G, Noordeloos ME, Santini A, Shouche YS, Bezerra JDP, Dima B, Guarnaccia V, Imrefi I, Jurjević Ž, Knapp DG, Kovács GM, Magistà D, Perrone G, Rämä T, Rebriev YA, Shivas RG, Singh SM, Souza-Motta CM, Thangavel R, Adhasure NN, Alexandrova AV, Alfenas AC, Alfenas RF, Alvarado P, Alves AL, Andrade DA, Andrade JP, Barbosa RN, Barili A, Barnes CW, Baseia IG, Bellanger JM, Berlanas C, Bessette AE, Bessette AR, Biketova AY, Bomfim FS, Brandrud TE, Bransgrove K, Brito ACQ, Cano-Lira JF, Cantillo T, Cavalcanti AD, Cheewangkoon R, Chikowski RS, Conforto C, Cordeiro TRL, Craine JD, Cruz R, Damm U, de Oliveira RJV, de Souza JT, de Souza HG, Dearnaley JDW, Dimitrov RA, Dovana F, Erhard A, Esteve-Raventós F, Félix

- CR, Ferisin G, Fernandes RA, Ferreira RJ, Ferro LO, Figueiredo CN, Frank JL, Freire KTLS, García D, Gené J, Gêsiorska A, Gibertoni TB, Gondra RAG, Gouliamova DE, Gramaje D, Guard F, Gusmão LFP, Haitook S, Hirooka Y, Houbraken J, Hubka V, Inamdar A, Iturriaga T, Iturrieta-González I, Jadan M, Jiang N, Justo A, Kachalkin AV, Kapitonov VI, Karadelev M, Karakehian J, Kasuya T, Kautmanová I, Kruse J, Kušan I, Kuznetsova TA, Landell MF, Larsson KH, Lee HB, Lima DX, Lira CRS, Machado AR, Madrid H, Magalhães OMC, Majerova H, Malysheva EF, Mapperson RR, Marbach PAS, Martín MP, Martín-Sanz A, Matočec N, McTaggart AR, Mello JF, Melo RFR, Mešić A, Michereff SJ, Miller AN, Minoshima A, Molinero-Ruiz L, Morozova OV, Mosoh D, Nabe M, Naik R, Nara K, Nascimento SS, Neves RP, Olariaga I, Oliveira RL, Oliveira TGL, Ono T, Ordoñez ME, Ottoni AM, Paiva LM, Pancorbo F, Pant B, Pawłowska J, Peterson SW, Raudabaugh DB, Rodríguez-Andrade E, Rubio E, Rusevska K, Santiago ALCMA, Santos ACS, Santos C, Sazanov NA, Shah S, Sharma J, Silva BDB, Siquier JL, Sonawane MS, Stchigel AM, Svetasheva T, Tamakeaw N, Telleria MT, Tiago PV, Tian CM, Tkalčec Z, Tomashevskaya MA, Truong HH, Vecherskii MV, Visagie CM, Vizzini A, Yilmaz N, Zmitrovich IV, Zvyagina EA, Boekhout T, Kehlet T, Læssøe T, Groenewald JZ (2019) Fungal Planet description sheets: 868–950. *Persoonia* 42: 291–473. <https://doi.org/10.3767/persoonia.2019.42.11>
- Crous PW, Wingfield MJ, Schumacher RK, Akulov A, Bulgakov TS, Carnegie AJ, Jurjević Ž, Decock C, Denman S, Lombard L, Lawrence DP, Stack AJ, Gordon TR, Bostock RM, Burgess T, Summerell BA, Taylor PWJ, Edwards J, Hou LW, Cai L, Rossman AY, Wöhner T, Allen WC, Castlebury LA, Visagie CM, Groenewald JZ (2020) New and Interesting Fungi. 3. *Fungal Systematics and Evolution* 6(1): 157–231. <https://doi.org/10.3114/fuse.2020.06.09>
- Crous PW, Osieck ER, Jurjević Ž, Boers J, van Iperen AL, Starink-Willemse M, Dima B, Balashov S, Bulgakov TS, Johnston PR, Morozova OV, Pinruan U, Sommai S, Alvarado P, Decock CA, Lebel T, McMullan-Fisher S, Moreno G, Shivas RG, Zhao L, Abdollahzadeh J, Abrinbana M, Ageev DV, Akhmetova G, Alexandrova AV, Altés A, Amaral AGG, Angelini C, Antonín V, Arenas F, Asselman P, Badali F, Baghela A, Bañares A, Barreto RW, Baseia IG, Bellanger JM, Berraf-Tebbal A, Biketova AY, Bukharova NV, Burgess TI, Cabero J, Câmara MPS, Cano-Lira JF, Ceryngier P, Chávez R, Cowan DA, de Lima AF, Oliveira RL, Denman S, Dang QN, Dovana F, Duarte IG, Eichmeier A, Erhard A, Esteve-Raventós F, Fellin A, Ferisin G, Ferreira RJ, Ferrer A, Finy P, Gaya E, Geering ADW, Gil-Durán C, Glässnerová K, Glushakova AM, Gramaje D, Guard FE, Guarnizo AL, Haelewaters D, Halling RE, Hill R, Hirooka Y, Hubka V, Iliushin VA, Ivanova DD, Ivanushkina NE, Jangsantear P, Justo A, Kachalkin AV, Kato S, Khamsuntorn P, Kirtsideli IY, Knapp DG, Kochkina GA, Koukol O, Kovács GM, Kruse J, Kumar TKA, Kušan I, Læssøe T, Larsson E, Lebeuf R, Levicán G, Loizides M, Marinho P, Luangsa-ard JJ, Lukina EG, Magaña-Dueñas V, Maggs-Köling G, Malysheva EF, Malysheva VF, Martín B, Martín MP, Matočec N, McTaggart AR, Mehrabi-Koushki M, Mešić A, Miller AN, Mironova P, Moreau P-A, Morte A, Müller K, Nagy LG, Nanu S, Navarro-Ródenas A, Nel WJ, Nguyen TH, Nóbrega TF, Noordeloos ME, Olariaga I, Overton BE, Ozerskaya SM, Palani P, Pancorbo F, Papp V, Pawłowska J, Pham TQ, Phosri C, Popov ES, Portugal A, Pošta A, Reschke K, Reul M, Ricci GM, Rodríguez A, Romanowski J, Ruchikachorn N, Saar I, Safi A, Sakolrak B, Salzmann F, Sandoval-Denis M, Sangwichein E, Sanhueza L, Sato T, Sastoque A, Senn-Irlet B, Shibata A, Siepe K, Somrithipol S, Spetik M, Sridhar P, Stchigel AM, Stusko-va K, Suwannasai N, Tan YP, Thangavel R, Tiago I, Tiwari S, Tkalčec Z, Tomashevskaya MA, Tonegawa C, Tran HX, Tran NT, Trovão J, Trubitsyn VE, Van Wyk J, Vieira WAS, Vila J, Visagie CM, Vizzini A, Volobuev SV, Vu DT, Wangsawat N, Yaguchi T, Ercole E, Ferreira BW, de Souza AP, Vieira BS, Groenewald JZ (2021) Fungal Planet description sheets: 1284–1382. *Persoonia* 47: 178–374. <https://doi.org/10.3767/persoonia.2021.47.06>

- Crous PW, Jurjević Z, Balashov S, De la Peña-Lastra S, Mateos A, Pinruan U, Rigueiro-Rodríguez A, Osieck ER, Altés A, Czachura P, Esteve-Raventós F, Gunaseelan S, Kaliyaperumal M, Larsson E, Luangsa-ard JJ, Moreno G, Pancorbo F, Piątek M, Sommai S, Somrithipol S, Asif M, Delgado G, Flakus A, Illescas T, Kezo K, Khamsuntorn P, Kubátová A, Labuda R, Lavoise C, Lebel T, Lueangjaroenkit P, Maciá-Vicente JG, Paz A, Saba M, Shivas RG, Tan YP, Wingfield MJ, Aas T, Abramczyk B, Ainsworth AM, Akulov A, Alvarado P, Armada F, Assyov B, Avchar R, Avesani M, Bezerra JL, Bhat JD, Bilański P, Bily DS, Boccardo F, Bozok F, Campos JC, Chaimongkol S, Chellappan N, Costa MM, Dalecká M, Darmostuk V, Daskalopoulos V, Dearnaley J, Dentinger BTM, De Silva NI, Dhotre D, Carlavilla JR, Doungsa-ard C, Dovana F, Erhard A, Ferro LO, Gallegos SC, Giles CE, Gore G, Gorfer M, Guard FE, Hanson SA, Haridev P, Jankowiak R, Jeffers SN, Kandemir H, Karich A, Kisło K, Kiss L, Krisai-Greilhuber I, Latha KPD, Lorenzini M, Lumyong S, Manimohan P, Manjón JL, Maula F, Mazur E, Mesquita NLS, Młynek K, Mongkolsamrit S, Morán P, Murugadoss R, Nagarajan M, Nalumpang S, Noisripoom W, Nosalj S, Novaes QS, Nowak M, Pawłowska J, Peiger M, Pereira OL, Pinto A, Plaza M, Polemis E, Polhorský A, Ramos DO, Raza M, Rivas-Ferreiro M, Rodriguez-Flakus P, Ruszkiewicz-Michalska M, Sánchez A, Santos A, Schüller A, Scott PA, Şen İ, Shelke D, Śliwa L, Solheim H, Sonawane H, Strašitáková D, Stryjak-Bogacka M, Sudsanguan M, Suwannarach N, Suz LM, Syme K, Taşkın H, Tennakoon DS, Tomka P, Vaghefi N, Vasan V, Vauras J, Wiktorowicz D, Villarreal M, Vizzini A, Wrzosek M, Yang X, Yingkunchao W, Zapparoli G, Zervakis GI, Groenewald JZ (2024) Fungal Planet description sheets: 1614–1696. *Fungal Systematics and Evolution* 13: 183–439. <https://doi.org/10.3114/fuse.2024.13.11>
- Damm U, Verkley GJM, Crous PW, Fourie PH, Haegi A, Riccioni L (2008) Novel *Paraconiothyrium* species on stone fruit trees and other woody hosts. *Persoonia* 20: 9–17. <https://doi.org/10.3767/003158508X286842>
- Das K, Lee SY, Choi HW, Eom AH, Cho YJ, Jung HY (2020) Taxonomy of *Arthrinium minutisporum* sp. nov., *Pezicula neosporulosa*, and *Acrocalymma pterocarpi*: New Records from Soil in Korea. *Mycobiology* 48: 450–463. <https://doi.org/10.1080/12298093.2020.1830741>
- Dong W, Wang B, Hyde KD, McKenzie EH, Raja HA, Tanaka K, Abdel-Wahab MA, Abdel-Aziz FA, Doilom M, Phookamsak R, Hongsanan S (2020) Freshwater Dothideomycetes. *Fungal Diversity* 105: 319–575. <https://doi.org/10.1007/s13225-020-00463-5>
- Ellis MB (1971) Dematiaceous Hyphomycetes. Commonwealth Mycological Institute. Kew, United Kingdom. <https://doi.org/10.1079/9780851986180.0000>
- Farr DF, Rossman AY (2025) Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/fungaldatabases> [accessed May 1, 2025]
- Farr DF, Miller ME, Bruton BD (1998) *Rhizopycnis vagum* gen. et sp. nov., a new coelomycetous fungus from roots of melons and sugarcane. *Mycologia* 90(2): 290–296. <https://doi.org/10.1080/00275514.1998.12026909>
- Fontecha G, Montes K, Ortiz B, Galindo C, Braham S (2019) Identification of cryptic species of four *Candida* complexes in a culture collection. *Journal of Fungi* (Basel, Switzerland) 5(4): 117. <https://doi.org/10.3390/jof5040117>
- Geethanjali P, Jayashankar M (2016) A review on litter decomposition by soil fungal community. *IOSR Journal of Pharmacy and Biological Sciences* 11(4): 1–3. <https://doi.org/10.9790/3008-1104030103>
- Giweta M (2020) Role of litter production and its decomposition, and factors affecting the processes in a tropical forest ecosystem: A review. *Journal of Ecology and Environment* 44: 1–9. <https://doi.org/10.1186/s41610-020-0151-2>

- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61(4): 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Gonçalves MF, Vicente TF, Esteves AC, Alves A (2019) *Neptunomyces aureus* gen. et sp. nov. (Didymosphaeriaceae, Pleosporales) isolated from algae in Ria de Aveiro, Portugal. *MycoKeys* 60: 31. <https://doi.org/10.3897/mycokeys.60.37931>
- Griffiths HM, Ashton LA, Parr CL, Eggleton P (2021) The impact of invertebrate decomposers on plants and soil. *The New Phytologist* 231(6): 2142–2149. <https://doi.org/10.1111/nph.17553>
- Guarnaccia V, Martino I, Brondino L, Gullino ML (2022) *Paraconiothyrium fuckelii*, *Diaporthe eres* and *Neocosmospora parceramosa* causing cane blight of red raspberry in Northern Italy. *Journal of Plant Pathology* 104(2): 683–698. <https://doi.org/10.1007/s42161-022-01068-4>
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hashimoto A, Matsumura M, Hirayama K, Fujimoto R, Tanaka K (2017) Pseudodidymellaceae fam. nov.: Phylogenetic affiliations of mycopappus-like genera in Dothideomycetes. *Studies in Mycology* 87(1): 187–206. <https://doi.org/10.1016/j.simyco.2017.07.002>
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Boonmee S, Lücking R, Pem D, Bhat DJ, Liu N, Tennakoon DS, Karunarathna A, Jiang SH, Jones EBG, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali D, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu YZ, Dissanayake AJ, Zeng XY, Luo Z, Tian Q, Phukhamsakda C, Thambugala KM, Dai D, Chethana KWT, Ertz D, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwanna C, Wijesinghe SN, Konta S, Niranjana M, Zhang SN, Ariyawansa HA, Jiang HB, Zhang JF, de Silva NI, Thiagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Aluthmuhandiram JVS, Abeywickrama PD, Bao DF, Devadatha B, Wu HX, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Chomnunti P, Samarakoon MC, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Bhunjun CS, Xu JC, Zheng J, Liu G, Feng Y, Xie N (2020) Refined families of Dothideomycetes: Dothideomycetidae and Pleosporomycetidae. *Mycosphere: Journal of Fungal Biology* 11(1): 1553–2107. <https://doi.org/10.5943/mycosphere/11/1/13>
- Htet ZH, Mapook A, Chethana KT (2024) Molecular taxonomy reveals new records of *Chromolaenicola* (Didymosphaeriaceae, Pleosporales) and potential antibacterial properties. *Studies in Fungi* 9(1): e006. <https://doi.org/10.48130/sif-0024-0006>
- Hyde KD, Jones EBG, Liu JK, Ariyawansa H, Boehm E, Boonmee S, Braun U, Chomnunti P, Crous PW, Dai DQ, Diederich P, Dissanayake A, Doilom M, Doveri F, Hongsanan S, Jayawardena R, Lawrey JD, Li YM, Liu YX, Lücking R, Monkai J, Muggia L, Nelsen MP, Pang KL, Phookamsak R, Senanayake I, Shearer CA, Suetrong S, Tanaka K, Thambugala KM, Wijayawardene NN, Wikee S, Wu HX, Zhang Y, Aguirre-Hudson B, Alias SA, Aptroot A, Bahkali AH, Bezerra JL, Bhat DJ, Camporesi E, Chukeatirote E, Gueidan C, Hawksworth DL, Hirayama K, Hoog SD, Kang JC, Knudsen K, Li WJ, Li XH, Liu ZY, Mapook A, McKenzie EHC, Miller AN, Mortimer PE, Phillips AJL, Raja HA, Scheuer C, Schumm F, Taylor JE, Tian Q, Tibpromma S, Wanasinghe DN, Wang Y, Xu JC, Yan JY, Yacharoen S, Zhang M (2013) Families of Dothideomycetes. *Fungal Diversity* 63: 1–313. <https://doi.org/10.1007/s13225-013-0263-4>
- Hyde KD, Hongsanan S, Jeewon R, Bhat DJ, McKenzie EHC, Jones EBG, Phookamsak R, Ariyawansa HA, Boonmee S, Zhao Q, Abdel-Aziz FA, Abdel-Wahab MA, Banmai S,

- Chomnunti P, Cui BK, Daranagama DA, Das K, Dayarathne MC, de Silva NI, Dissanayake AJ, Doilom M, Ekanayaka AH, Gibertoni TB, Go'és-Neto A, Huang SK, Jayasiri SC, Jayawardena RS, Konta S, Lee HB, Li WJ, Lin CG, Liu JK, Lu YZ, Luo ZL, Manawasinghe IS, Manimohan P, Mapook A, Niskanen T, Norphanphoun C, Papizadeh M, Perera RH, Phukhamsakda C, Richter C, de Santiago ALCMA, Drechsler-Santos ER, Senanayake IC, Tanaka K, Tennakoon TMDS, Thambugala KM, Tian Q, Tibpromma S, Thongbai B, Vizzini A, Wanasinghe DN, Wijayawardene NN, Wu HX, Yang J, Zeng XY, Zhang H, Zhang JF, Bulgakov TS, Camporesi E, Bahkali AH, Amoozegar AM, Araujo-Neta LS, Ammirati JF, Baghela A, Bhatt RP, Bojantchev S, Buyck B, da Silva GA, de Lima CLF, de Oliveira RJV, de Souza CAF, Dai YC, Dima B, Duong TT, Ercole E, Mafalda-Freire F, Ghosh A, Hashimoto A, Kamolhan S, Kang JC, Karunarathna SC, Kirk PM, Kytovuori I, Lantieri A, Liimatainen K, Liu ZY, Liu XZ, Lu'cking R, Medardi G, Mortimer PE, Nguyen TTT, Promputtha I, Raj KNA, Reck MA, Lumyong S, Shahzadeh-Fazeli SA, Stadler M, Soudi MR, Su HY, Takahashi T, Tangthirasun N, Uniyal P, Wang Y, Wen TC, Xu JC, Zhang ZK, Zhao YC, Zhou JZ, Zhu L (2016) Fungal diversity notes 367–490: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 80: 1–270. <https://doi.org/10.1007/s13225-016-0373-x>
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Chethana KWT, Clericuzio M, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, He MQ, Hongsan S, Huang SK, Jayasiri SC, Jayawardena RS, Karunarathna A, Konta S, Kusan I, Lee H, Li J, Lin CG, Liu NG, Lu YZ, Luo ZL, Manawasinghe IS, Mapook A, Perera RH, Phookamsak R, Phukhamsakda C, Siedlecki I, Soares AM, Tennakoon DS, Tian Q, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Abdel-Aziz FA, Li WJ, Senanayake IC, Shang QJ, Daranagama DA, De Silva NI, Thambugala KM, Abdel-Wahab MA, Bahkali AH, Berbee ML, Boonmee S, Bhat DJ, Bulgakov TS, Buyck B, Camporesi E, Castañeda-Ruiz RF, Chomnunti P, Doilom M, Dovana F, Gibertoni TB, Jadan M, Jeewon R, Jones GEB, Kang JC, Karunarathna SC, Lim YW, Liu JK, Liu ZY, Plautz Jr HL, Lumyong S, Maharachchikumbura SSN, Matočec N, Mckenzie EHC, Mešić A, Miller D, Pawłowska J, Pereira OL, Promputtha I, Romero AI, Ryvarden L, Su HY, Suetrong S, Tkálčec Z, Vizzini A, Wen TC, Wisitrassameewong K, Wrzosek M, Xu JC, Zhao Q, Zhao RL, Mortimer PE (2017) Fungal diversity notes 603–708: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 87: 1–235. <https://doi.org/10.1007/s13225-017-0391-3>
- Hyde KD, Dong Y, Phookamsak R, Jeewon R, Bhat DJ, Jones EBG, Liu NG, Abeywickrama PD, Mapook A, Wei D, Perera RH, Manawasinghe IS, Pem D, Bundhun D, Karunarathna A, Ekanayaka AH, Bao DF, Li J, Samarakoon MC, Chaiwan N, Lin CG, Phutthacharoen K, Zhang SN, Senanayake IC, Goonasekara ID, Thambugala KM, Phukhamsakda C, Tennakoon DS, Jiang HB, Yang J, Zeng M, Huanraluek N, Liu JK, Wijesinghe SN, Tian Q, Tibpromma S, Brahmanage RS, Boonmee S, Huang SK, Thiyagaraja V, Lu YZ, Jayawardena RS, Dong W, Yang EF, Singh SK, Singh SM, Rana S, Lad SS, Anand G, Devadatha B, Niranjana M, Sarma VV, Liimatainen K, Aguirre-Hudson B, Niskanen T, Overall A, Alvarenga RLM, Gibertoni TB, Pfliegler WP, Horváth E, Imre A, Alves AL, da Silva Santos AC, Tiago PV, Bulgakov TS, Wanasinghe DN, Bahkali AH, Doilom M, Elgorban AM, Maharachchikumbura SSN, Rajeshkumar KC, Haelewaters D, Mortimer PE, Zhao Q, Lumyong S, Xu J, Sheng J (2020) Fungal diversity notes 1151–1276: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 100(1): 5–277. <https://doi.org/10.1007/s13225-020-00439-5>
- Index Fungorum (2025) Index Fungorum. <http://www.indexfungorum.org/names/Names.asp> [accessed 6 May 2025]

- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat DJ, Buyck B, Cai L, Dai YC, Abd–Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa–ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad–Nejhad M, Nilsson H, Pang KA, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel–Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The faces of fungi database: Fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jayasiri SC, Hyde KD, Jones EBG, McKenzie EHC, Jeewon R, Phillips AJL, Bhat DJ, Wanasinghe DN, Liu JK, Lu YZ, Kang JC, Xu J, Karunarathna SC (2019) Diversity, morphology and molecular phylogeny of Dothideomycetes on decaying wild seed pods and fruits. *Mycosphere: Journal of Fungal Biology* 10: 1–186. <https://doi.org/10.5943/mycosphere/10/1/1>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Khan AL, Hamayun M, Hussain J, Kang SM, Lee IJ (2012) The newly isolated endophytic fungus *Paraconiothyrium* sp. LK1 produces ascotoxin. *Molecules (Basel, Switzerland)* 17(1): 1103–1112. <https://doi.org/10.3390/molecules17011103>
- Konta S, Tibpromma S, Karunarathna SC, Samarakoon MC, Steven LS, Mapook A, Boonmee S, Senwanna C, Balasuriya A, Eungwanichayapant PD, Hyde KD (2023) Morphology and multigene phylogeny reveal ten novel taxa in Ascomycota from terrestrial palm substrates (Arecaceae) in Thailand. *Mycosphere: Journal of Fungal Biology* 14(1): 107–152. <https://doi.org/10.5943/mycosphere/14/1/2>
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SSN, McKenzie EHC, Phookamsak R, Phukhamsakda C, Shenoy BD, Abdel–Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, Dsouza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsan S, Jayasiri SC, Jayawardena RS, Karunarathna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Peršoh D, Pinruan U, Senanayake IC, Somrithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Wisitrassameewong K, Zeng XY, Abdel–Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Lewis J, Li XH, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sri–Indrasutdhi V, Tian CM, Verbeken A, von Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E (2015) Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72(1): 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Liu S, He G, Fang H, Xu S, Bai S (2022) Effects of dissolved oxygen on the decomposers and decomposition of plant litter in lake ecosystem. *Journal of Cleaner Production* 372: 133837. <https://doi.org/10.1016/j.jclepro.2022.133837>
- Lu B, Hyde KD, Ho WH, Tsui KM, Taylor JE, Wong KM, Yanna M, Zhou DQ (2000) Checklist of Hong Kong Fungi. *Fungal Diversity Research Series*, Hong Kong.
- Lücking R, Aime MC, Robbertse B, Miller AN, Aoki T, Ariyawansa HA, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, Hawksworth DL (2021) Fungal taxonomy and sequence-based nomenclature. *Nature Microbiology* 6(5): 540–548. <https://doi.org/10.1038/s41564-021-00888-x>

- Manish T, Gupta RC, Yogesh J (2014) *Spegazzinia tessarthra* isolated as a true endophyte from lichen *Heterodermia flabellate*. Indian Phytopathology 67: 109–110.
- Manoharachary C, Kunwar IK (2010) *Spegazzinia* species from India. Taxonomy and Ecology of Indian fungi. IK Internat Pvt, New Delhi.
- Matsushima T (1980) Matsushima Mycological Memoirs No. 1. Saprophytic Microfungi from Taiwan, Part 1. Hyphomycetes. Matsushima Fungus Collection, Kobe, 82.
- Mena-Portales J, Cantillo-Perez T, Minter DW (2017) A new species of the conidial fungal genus *Spegazzinia* (Pleosporales Didymosphaeriaceae) collected on sugarcane in Cuba. Phytotaxa 331(2): 295–298. <https://doi.org/10.11646/phytotaxa.331.2.14>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees. SC10 Workshop on Gateway Computing Environments (GCE10). <https://doi.org/10.1109/GCE.2010.5676129>
- Mortimer PE, Jeewon R, Xu JC, Lumyong S, Wanasinghe DN (2021) Morpho-phylo taxonomy of novel Dothideomycetous fungi associated with dead woody twigs in Yunnan Province, China. Frontiers in Microbiology 12: 654683. <https://doi.org/10.3389/fmicb.2021.654683>
- Munk A (1956) On *Metasphaeria coccodes* (Karst.) Sacc. (Massarinaceae n. fam.). Friesia 5: 305.
- Nagraj T (1978) Genera Coelomycetum. 16. *Fibulocoela* form-gen nov, A Coelomycete with basidiomycetous affinities. Canadian Journal of Botany 56: 1485–1491. <https://doi.org/10.1139/b78-173>
- Nakashima KI, Tomida J, Hirai T, Kawamura Y, Inoue M (2019) Paraconiothins A–J: Sesquiterpenoids from the endophytic fungus *Paraconiothyrium brasiliense* ECN258. Journal of Natural Products 82: 3347–3356. <https://doi.org/10.1021/acs.jnatprod.9b00638>
- Nylander JAA (2004) MrModeltest 2.0. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Pascoal C, Fernandes I, Seena S, Danger M, Ferreira V, Cássio F (2021) Linking Microbial Decomposer Diversity to Plant Litter Decomposition and Associated Processes in Streams. In: Swan CM, Boyero L, Canhoto C (Eds) The Ecology of Plant Litter Decomposition in Stream Ecosystems. Springer, Cham. https://doi.org/10.1007/978-3-030-72854-0_9
- Paul NC, Lee HB (2014) First record of endophytic *Paraconiothyrium brasiliense* isolated from Chinese maple leaves in Korea. Korean Journal of Mycology 42: 349–352. <https://doi.org/10.4489/KJM.2014.42.4.349>
- Pei Z, Leppert KN, Eichenberg D, Bruehlheide H, Niklaus PA, Buscot F, Gutknecht JLM (2017) Leaf litter diversity alters microbial activity, microbial abundances, and nutrient cycling in a subtropical forest ecosystem. Biogeochemistry 134: 163–181. <https://doi.org/10.1007/s10533-017-0353-6>
- Phookamsak R, Hyde KD, Jeewon R, Bhat DJ, Jones EBG, Maharachchikumbura SSN, Raspe O, Karunarathna SC, Wanasinghe DN, Hongsan S, Doilom M, Tennakoon DS, Machado AR, Firmino AL, Ghosh A, Karunarathna A, Mešić A, Dutta AK, Thongbai B, Devadatha B, Norphanphoun C, Senwanna C, Wei D, Pem D, Ackah FK, Wang G-N, Jiang H-B, Madrid H, Lee HB, Goonasekara ID, Manawasinghe IS, Kušan I, Cano J, Gené J, Li J, Das K, Acharya K, Raj KNA, Latha KPD, Chethana KWT, He M-Q, Dueñas M, Jadan M, Martín MP, Samarakoon MC, Dayarathne MC, Raza M, Park MS, Telleria MT, Chaiwan N, Matočec N, de Silva NI, Pereira OL, Singh PN, Manimohan P, Uniyal P, Shang Q-J, Bhatt RP, Perera RH, Alvarenga RLM, Nogal-Prata S, Singh SK, Vadthananarat S, Oh S-Y, Huang S-K, Rana S, Konta S, Paloi S, Jayasiri SC, Jeon SJ, Mehmood T, Gibertoni TB, Nguyen TTT, Singh U, Thiyagaraja V, Sarma VV, Dong W, Yu X-D, Lu Y-Z, Lim YW, Chen Y, Tkáčec Z, Zhang Z-F, Luo Z-L, Daranagama DA, Thambugala KM,

- Tibpromma S, Camporesi E, Bulgakov TS, Dissanayake AJ, Senanayake IC, Dai DQ, Tang L-Z, Khan S, Zhang H, Promputtha I, Cai L, Chomnunti P, Zhao R-L, Lumyong S, Boonmee S, Wen T-C, Mortimer PE, Xu J (2019) Fungal diversity notes 929–1035, taxonomic and phylogenetic contributions on genera and species of fungi. *Fungal Diversity* 95(1): 1–273. <https://doi.org/10.1007/s13225-019-00421-w>
- Rambaut A (2012) FigTree Version 1.4.0. Edinburgh: Institute of Evolutionary Biology; University of Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/> [accessed May 10, 2025]
- Rathnayaka AR, Chethana KWT, Manawasinghe IS, Wijesinghe SN, de Silva NI, Tennakoon DS, Phillips AJL, Liu JK, Jones EBG, Wang Y, Hyde KD (2023) *Lasiodiplodia*: Generic revision by providing molecular markers, geographical distribution and haplotype diversity. *Mycosphere: Journal of Fungal Biology* 14(1): 1254–1339. <https://doi.org/10.5943/mycosphere/14/1/14>
- Rathnayaka AR, Tennakoon DS, Jones GE, Wanasinghe DN, Bhat DJ, Priyashantha AH, Stephenson SL, Tibpromma S, Karunarathna SC (2025) Significance of precise documentation of hosts and geospatial data of fungal collections, with an emphasis on plant-associated fungi. *New Zealand Journal of Botany* 63(2–3): 462–489. <https://doi.org/10.1080/0028825X.2024.2381734>
- Rehner S (2001) Primers for Elongation Factor 1- α (EF1- α). <http://ocid.NACSE.ORG/research/deephyphae/EF1primer.pdf>
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Saccardo PA (1880) Fungi *Gallici lecti* a cl. viris P. Brunaud, Abb. Letendre, A. Malbranche, J. Therry vel editi in *Mycotheca Gallica* C. Roumeguèri. Series II. *Michelia* 2: 39–135.
- Samaradiwakara NP, Zou X, Yang Y, de Farias ARG, Gajanayake AJ, Tennakoon DS, Manawasinghe IS, Dong Z, Luo M, Lumyong S (2022) First report of *Pseudopithomyces maydicus* causing leaf spots in strawberry. *Crop Protection (Guildford, Surrey)* 162: 106081. <https://doi.org/10.1016/j.cropro.2022.106081>
- Samarakoon BC, Phookamsak R, Wanasinghe DN, Chomnunti P, Hyde KD, McKenzie EH, Promputtha I, Xu JC, Li YJ (2020) Taxonomy and phylogenetic appraisal of *Spegazzinia musae* sp. nov. and *S. deightonii* (Didymosphaeriaceae, Pleosporales) on Musaceae from Thailand. *Mycoskeys* 70: 19. <https://doi.org/10.3897/mycokeys.70.52043>
- Sayer EJ (2006) Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. *Biological Reviews of the Cambridge Philosophical Society* 81: 1–31. <https://doi.org/10.1017/S1464793105006846>
- Schroeter SA, Eveillard D, Chaffron S, Zoppi J, Kampe B, Lohmann P, Jehmlich N, von Bergen M, Sanchez-Arcos C, Pohnert G, Taubert M (2022) Microbial community functioning during plant litter decomposition. *Scientific Reports* 12(1): 7451. <https://doi.org/10.1038/s41598-022-11485-1>
- Senanayake IC, Pem D, Rathnayaka AR, Wijesinghe SN, Tibpromma S, Wanasinghe DN, Phookamsak R, Kularathnage ND, Gomdola D, Harishchandra D, Dissanayake LS, Xiang M, Ekanayaka AH, McKenzie EHC, Hyde KD, Zhang H, Xie N (2022) Predicting global numbers of teleomorphic ascomycetes. *Fungal Diversity* 114(1): 237–278. <https://doi.org/10.1007/s13225-022-00498-w>
- Shoemaker RA, Babcock CE, Irwin JAG (1991) *Massarina walkeri* n. sp., the teleomorph of *Acrocalymma medicaginis* from *Medicago sativa* contrasted with *Leptosphaeria pratensis*, *L. weimeri* n. sp., and *L. viridella*. *Canadian Journal of Botany* 69: 569–573. <https://doi.org/10.1139/b91-077>

- Species Fungorum (2025) Species Fungorum. <http://www.indexfungorum.org> [accessed on 15 April 2025]
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* (Oxford, England) 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Sutton BC (1980) The coelomycetes. Fungi Imperfecti with Pycnidia, Acervuli and Stromata. Commonwealth Mycological Institute.
- Tanaka K, Harada Y, Barr ME (2005) Bambusicolous fungi in Japan (3): A new combination, *Kalmusia scabrispora*. *Mycoscience* 46: 110–113. <https://doi.org/10.1007/S10267-004-0224-6>
- Tanaka K, Hirayama K, Yonezawa H, Sato G, Toriyabe A, Kudo H, Hashimoto A, Matsumura M, Harada Y, Kurihara Y, Shirouzu T (2015) Revision of the Massarineae (Pleosporales, Dothideomycetes). *Studies in Mycology* 82: 75–136. <https://doi.org/10.1016/j.simyco.2015.10.002>
- Tedersoo L, Anslan S, Bahram M, Kõljalg U, Abarenkov K (2020) Identifying the ‘unidentified’ fungi: A global-scale long-read third-generation sequencing approach. *Fungal Diversity* 103: 273–293. <https://doi.org/10.1007/s13225-020-00456-4>
- Tennakoon DS, Jeewon R, Kuo CH, Hyde KD (2018) Phylogenetic and morphological characterization of *Byssosphaeria macarangae* sp. nov., and *B. taiwanense* sp. nov. from *Macaranga tanarius*. *Phytotaxa* 364(3): 211–226. <https://doi.org/10.11646/phytotaxa.364.3.1>
- Tennakoon DS, Thambugala KM, Wanasinghe DN, Gentekaki E, Promputtha I, Kuo CH, Hyde KD (2020) Additions to Phaeosphaeriaceae (Pleosporales): *Elongaticollum* gen. nov., *Ophiosphaerella taiwanensis* sp. nov., *Phaeosphaeriopsis beaucarneae* sp. nov. and a new host record of *Neosetophoma poaceicola* from Musaceae. *MycKeys* 70: 59. <https://doi.org/10.3897/mycokeys.70.53674>
- Tennakoon DS, Kuo CH, Maharachchikumbura SSN, Thambugala KM, Gentekaki E, Phillips AJ, Bhat DJ, Wanasinghe DN, de Silva NI, Promputtha I, Hyde KD (2021) Taxonomic and phylogenetic contributions to *Celtis formosana*, *Ficus ampelas*, *F. septica*, *Macaranga tanarius* and *Morus australis* leaf litter inhabiting microfungi. *Fungal Diversity* 108(1): 1–215. <https://doi.org/10.1007/s13225-021-00474-w>
- Tennakoon DS, Thambugala KM, Silva NID, Suwannarach N, Lumyong S (2022) A taxonomic assessment of novel and remarkable fungal species in Didymosphaeriaceae (Pleosporales, Dothideomycetes) from plant litter. *Frontiers in Microbiology* 13: e1016285. <https://doi.org/10.3389/fmicb.2022.1016285>
- Tennakoon DS, de Silva NI, Maharachchikumbura SS, Bhat DJ, Kumla J, Suwannarach N, Lumyong S (2023) Exploring More on Dictyosporiaceae: The Species Geographical Distribution and Intriguing Novel Additions from Plant Litter. *Diversity* 15(3): e410. <https://doi.org/10.3390/d15030410>
- Tennakoon DS, Thambugala KM, de Silva NI, Song HY, Suwannarach N, Chen FS, Hu DM (2024) An overview of Melanommataceae (Pleosporales, Dothideomycetes): Current insight into the host associations and geographical distribution with some interesting novel additions from plant litter. *MycKeys* 106: 43. <https://doi.org/10.3897/mycokeys.106.125044>
- Thambugala KM, Wanasinghe DN, Phillips AJL, Camporesi E, Bulgakov TS, Phukham-sakda C, Ariyawansa HA, Goonasekara ID, Phookamsak R, Dissanayake A, Tennakoon DS (2017) Mycosphere notes 1–50: Grass (Poaceae) inhabiting Dothideomycetes. *Mycosphere: Journal of Fungi* 8(4): 697–796. <https://doi.org/10.5943/mycosphere/8/4/13>

- Tianyu Z (2009). Flora Fungorum Sincorum (Vol 31): 26 Genera of Dematiaceous Dityosporous Hyphomycetes Excluding Alternaria. Science Press, Beijing.
- Trakunyingcharoen T, Lombard L, Groenewald JZ, Cheewangkoon R, Toanun C, Alfenas AC, Crous PW (2014) Mycoparasitic species of *Sphaerellopsis*, and allied lichenicolous and other genera. IMA Fungus 5: 391–414. <https://doi.org/10.5598/ima fungus.2014.05.02.05>
- Verkley GJ, da Silva M, Wicklow DT, Crous PW (2004) *Paraconiothyrium*, a new genus to accommodate the mycoparasite *Coniothyrium minitans*, anamorphs of *Paraphaeosphaeria*, and four new species. Studies in Mycology 50: 323–335.
- Verkley GJM, Dukik K, Renfurm R, Göker M, Stielow JB (2014) Novel genera and species of coniothyrium-like fungi in Montagnulaceae (Ascomycota). Persoonia 32: 25–51. <https://doi.org/10.3767/003158514X679191>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wanasinghe DN, Jones EBG, Camporesi E, Dissanayake AJ, Kamolhan S, Mortimer PE, Xu J, Abd-Elsalam KA, Hyde KD (2016) Taxonomy and phylogeny of *Laburnicola* gen. nov. and *Paramassariosphaeria* gen. nov. (Didymosphaeriaceae, Massariaceae, Pleosporales). Fungal Biology 120(11): 1354–1373. <https://doi.org/10.1016/j.funbio.2016.06.006>
- Wanasinghe DN, Ren GC, Xu JC, Cheewangkoon R, Mortimer PE (2022) Insight into the taxonomic resolution of the pleosporalean species associated with dead woody litter in natural forests from Yunnan, China. Journal of Fungi (Basel, Switzerland) 8(4): 375. <https://doi.org/10.3390/jof8040375>
- Wang J, Shao S, Liu C, Song Z, Liu S, Wu S (2021) The genus *Paraconiothyrium*: Species concepts, biological functions, and secondary metabolites. Critical Reviews in Microbiology 47(6): 781–810. <https://doi.org/10.1080/1040841X.2021.1933898>
- Wardle DA, Bonner KI, Barker GM (2002) Linkages between plant litter decomposition, litter quality, and vegetation responses to herbivores. Functional Ecology 16(5): 585–595. <https://doi.org/10.1046/j.1365-2435.2002.00659.x>
- White TJ, Bruns T, Lee SJWT, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiagaraja V, Tibpromma S, Tkálčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa (2021). Mycosphere: Journal of Fungal Biology 13: 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Wu X, Tang X, Hu S, Zhou Z, Zhu W, Li H, Li Z, Ding H (2023) First report of *Pseudopithomyces chartarum* causing leaf spot on *Tetrapanax papyrifer* in Guizhou, China. Plant Disease 107(12): 4021. <https://doi.org/10.1094/PDIS-03-23-0453-PDN>
- Yasanthika W, Chethana K, Wanasinghe D, Tennakoon D, Al-Otibi F, Hyde K (2024) Taxonomic novelties of soil-inhabiting pleosporalean fungi in Thailand using baiting method. Phytotaxa 676(1): 1–24. <https://doi.org/10.11646/phytotaxa.676.1.1>

- Yuan Z, Druzhinina IS, Wang X, Zhang X, Peng L, Labbé J (2020) Insight into a highly polymorphic endophyte isolated from the roots of the halophytic seepweed *Suaeda salsa*: *Laburnicola rhizohalophila* sp. nov. (Didymosphaeriaceae, Pleosporales). *Fungal Biology* 124(5): 327–337. <https://doi.org/10.1016/j.funbio.2019.10.001>
- Zeng YL, Lu XJ, Wang MR, Zhu JN, Li QX, Zhu QP, Wang L, Bai ZF, Liu XH, Su ZZ, Lin FC (2023) Endophytic *Acrocalymma vagum* drives the microbiome variations for rice health. <https://doi.org/10.21203/rs.3.rs-2897955/v1>
- Zhang Y, Fournier J, Crous PW, Pointing SB, Hyde KD (2009) Phylogenetic and morphological assessment of two new species of *Amniculicola* and their allies (Pleosporales). *Persoonia* 23: 48–54. <https://doi.org/10.3767/003158509X472187>
- Zhang H, Hyde KD, Mckenzie EH, Bahkali AH, Zhou D (2012) Sequence data reveal phylogenetic affinities of *Acrocalymma aquatica* sp. nov., *Aquasubmersa mircensis* gen. et sp. nov. and *Clohesyomyces aquaticus* (freshwater coelomycetes). *Cryptogamie. Mycologie* 33: 333–346. <https://doi.org/10.7872/crym.v33.iss3.2012.333>
- Zhou X, Dong K, Tang Y, Huang H, Peng G, Wang D (2023) Research progress on the decomposition process of plant litter in wetlands: A review. *Water (Basel)* 15(18): 3246. <https://doi.org/10.3390/w15183246>